arXiv:1608.02091v1 [math.PR] 6 Aug 2016

Second-order asymptotics on distributions of maxima of bivariate elliptical arrays

Xin Liao[a] Zhichao Weng[b] Zuoxiang Peng[c*]

[a]Business School, University of Shanghai for Science and Technology, 200093 Shanghai, China

[b]School of Economics and Management, Fuzhou University, 350116 Fujian, China

[c]School of Mathematics and Statistics, Southwest University, 400715 Chongqing, China

Abstract Let $\{(\xi_{ni}, \eta_{ni}), 1 \leq i \leq n, n \geq 1\}$ be a triangular array of independent bivariate elliptical random vectors with the same distribution function as $(S_1, \rho_n S_1 + \sqrt{1-\rho_n^2} S_2)$, $\rho_n \in (0,1)$, where (S_1, S_2) is a bivariate spherical random vector. For the distribution function of radius $\sqrt{S_1^2 + S_2^2}$ belonging to the max-domain of attraction of the Weibull distribution, Hashorva (2006) derived the limiting distribution of maximum of this triangular array if convergence rate of ρ_n to 1 is given. In this paper, under the refinement of the rate of convergence of ρ_n to 1 and the second-order regular variation of the distributional tail of radius, precise second-order distributional expansions of the normalized maxima of bivariate elliptical triangular arrays are established.

Key words Bivariate elliptical triangular array; Maximum; Second-order expansion; Second-order regular variation.

AMS 2000 subject classification Primary 62E20, 60G70; Secondary 60F15, 60F05.

*Corresponding author. Email: pzx@swu.edu.cn

1 Introduction

Let $\{(\xi_{ni}, \eta_{ni}), 1 \le i \le n, n \ge 1\}$ be a triangular array of independent bivariate elliptical random vectors with stochastic representation

$$(\xi_{ni}, \eta_{ni}) \stackrel{d}{=} \left(S_1, \rho_n S_1 + \sqrt{1-\rho_n^2} S_2\right), 1 \le i \le n, n \ge 1, \tag{1.1}$$

where $\rho_n \in [-1, 1]$ and (S_1, S_2) be a bivariate spherical random vector with radius $R = \sqrt{S_1^2 + S_2^2}$. Here $\stackrel{d}{=}$ means equality in distribution. The bivariate maxima $\mathbf{M}_n$ is defined componentwise by

$$\mathbf{M}_n = (M_{n1}, M_{n2}) = \left(\max_{1\le i\le n} \xi_{ni}, \max_{1\le i\le n} \eta_{ni}\right).$$

If S_1 and S_2 are two independent standard Gaussian random variables, (1.1) just represents the well-known bivariate Gaussian triangular arrays studied by Hüsler and Reiss (1989) and Kabluchko et al. (2009) who showed that

$$\lim_{n\to\infty} \sup_{x,y\in\mathbb{R}} \left|\mathbb{P}\left(M_{n1} \le \bar{a}_n x + \bar{b}_n, M_{n2} \le \bar{a}_n y + \bar{b}_n\right) - K_\lambda(x, y)\right| = 0$$

if and only if the following Hüsler-Reiss condition

$$\lim_{n\to\infty} \frac{\bar{b}_n(1-\rho_n)}{\bar{a}_n} = 2\lambda^2 \in [0, \infty]$$

holds, where $K_\lambda(x, y)$ is the Hüsler-Reiss max-stable distribution, and the norming constants $\bar{b}_n$ satisfies $\sqrt{2\pi} n^{-1} \bar{b}_n \exp(\bar{b}_n^2/2) = 1$ and $\bar{a}_n = \bar{b}_n^{-1}$.

Motivated by the seminal work of Hüsler and Reiss (1989), Hashorva (2006) considered limit laws of maxima of bivariate elliptical triangular array with distribution function (df) F of radius R belonging to the max-domain of attraction of the Weibull extreme value distribution Ψ_α, $\alpha > 0$. Precisely, if the following condition

$$\lim_{n\to\infty} \frac{1-\rho_n}{a_n} = 2\lambda^2 \in (0, \infty) \tag{1.2}$$

holds with $a_n = 1 - G^{\leftarrow}\left(1 - n^{-1}\right)$, Hashorva (2006) showed that

$$\lim_{n\to\infty} \sup_{(x,y)\in(-\infty,0)^2} \left|\mathbb{P}\{M_{n1} \le 1 + a_n x, M_{n2} \le 1 + a_n y\} - H_{\alpha+\frac{1}{2},\lambda}(x,y)\right| = 0, \tag{1.3}$$

where G is the df of S_1 with upper endpoint 1, $G^{\leftarrow}$ is its generalised inverse function, and

$$H_{\alpha+\frac{1}{2},\lambda}(x,y) = \exp\left(-|x|^{\alpha+\frac{1}{2}}\psi_\alpha\left(\frac{1}{\sqrt{2|x|}}\left(\lambda + \frac{y-x}{2\lambda}\right)\right) - |y|^{\alpha+\frac{1}{2}}\psi_\alpha\left(\frac{1}{\sqrt{2|y|}}\left(\lambda + \frac{x-y}{2\lambda}\right)\right)\right).$$

Here, ψ_α is a df defined on $[-1,1]$ given by

$$\psi_\alpha(z) = \frac{\Gamma(\alpha+\frac{3}{2})}{\Gamma(\alpha+1)\sqrt{\pi}}\int_{-1}^{z}(1-s^2)^\alpha ds,\ \alpha > 0,\ z \in [-1,1],$$

where $\Gamma(\cdot)$ is Gamma function. For more work on extremal behaviors of extended Hüsler-Reiss bivariate Gaussian settings, see, e.g., Hooghiemstra and Hüsler (1996), Hashorva (2005, 2008), Frick and Reiss (2010), Hashorva et al. (2012), Hashorva and Weng (2013), Hashorva and Ling (2016), and Liao et al. (2016). For multivariate dependent Gaussian case, see Hashorva et al. (2015).

Recently, the studies on convergence rate of normalized maxima of Hüsler-Reiss bivariate Gaussian settings and their extensions have been received more attentions. For bivariate Gaussian triangular arrays, Hashorva et al. (2016) established the higher-order distributional expansions of normalized maxima under refined Hüsler-Reiss condition, and the uniform convergence rates of normalized maxima and the second-order expansion of the joint distributions of normalized maxima and minima were investigated by Liao and Peng (2014, 2015), respectively. For copula version of bivariate Gaussian triangular arrays, Frick and Reiss (2013) considered the penultimate and ultimate convergence rate for distributions of $(n(\max_{1\le i\le n}\Phi(\xi_{ni}) - 1), n(\max_{1\le i\le n}\Phi(\eta_{ni}) - 1))$, where $\Phi(\cdot)$ is the standard Gaussian distribution.

In this paper, we are interested in the second-order distributional expansions of bivariate normalized maxima as the distribution F of radius R belongs to the max-domain of attraction of Ψ_α,

$\alpha > 0$. From Resnick (1987), it follows that F is in the Weibull max-domain of attraction, i.e.,

$$\lim_{n\to\infty} \sup_{x\in\mathbb{R}} |F^n(\tilde{a}_n x + \tilde{b}_n) - \Psi_\alpha(x)| = 0$$

if and only if the upper endpoint $\omega := \sup\{t : F(t) < 1\} < \infty$, and $1 - F(\omega - s^{-1}) \in RV_{-\alpha}$, regularly varying functions with exponent $-\alpha$. Furthermore, the constants can be chosen as

$$\tilde{a}_n := \omega - F^{\leftarrow}(1 - n^{-1}),\ \tilde{b}_n := \omega,\ n > 1.$$

In order to get the desired results, we assume that $1 - F(\omega - s^{-1})$ has the properties of second-order regularly variation with the first-order parameter $-\alpha$ and the second-order parameter $\tau \le 0$ (written as $1 - F(\omega - s^{-1}) \in 2RV_{-\alpha,\tau}$), i.e., there exists some ultimately positive or negative function $A(t)$ with $\lim_{t\to\infty} A(t) = 0$ such that

$$\lim_{t\to\infty} \frac{\frac{1-F(\omega-(ts)^{-1})}{1-F(\omega-t^{-1})} - s^{-\alpha}}{A(t)} = s^{-\alpha}\frac{s^\tau - 1}{\tau}, \quad s > 0,$$

where $\frac{s^\tau-1}{\tau}$ is interpreted as $\log s$ when $\tau = 0$, c.f., de Haan and Ferreira (2006). Furthermore, we need the rate of convergence imposed on (1.2) as $\lambda \in (0, \infty)$. Assume that throughout this paper that there exists c_n with $\lim_{n\to\infty} c_n = 0$ such that

$$\lim_{n\to\infty} \frac{\lambda_n - \lambda}{c_n} = \gamma \in \mathbb{R} \tag{1.4}$$

holds with $\lambda_n = \left(\frac{1-\rho_n}{2a_n}\right)^{1/2}$ and $\lambda \in (0, \infty)$. For two extreme cases $\lambda = 0$ and $\lambda = \infty$, the analysis will be discussed with some other additional conditions related to ρ_n.

The rest of this paper is organized as follows. In section 2, we provide the main results and an example is illustrated to support our findings. Some auxiliary lemmas are given in Section 3, and all proofs of the main results are deferred to Section 4.

2 Main results

In this section, we provide the main results with respect to the second-order expansions on distributions of normalized maxima by assuming that $1-F(\omega-s^{-1}) \in 2RV_{-\alpha,\tau}$, $\alpha>0, \tau \le 0$. There are three cases to be considered, i.e., $\lambda \in (0,\infty)$, $\lambda=0$ and $\lambda=\infty$, respectively. For simplicity, we assume that the upper endpoint ω of F is 1, which implies that the upper endpoint of G is also 1. Throughout this paper, let a_n be the normalized constant satisfied $a_n = 1-G^{\leftarrow}(1-n^{-1})$. For the case of $\lambda \in (0,\infty)$, we need a second-order condition (1.4) refining the convergence rate of condition (1.2), and investigate in turn the following three cases: $\lambda^2+x+y+\frac{(x-y)^2}{4\lambda^2}<0$, $\lambda^2+x+y+\frac{(x-y)^2}{4\lambda^2}=0$ and $\lambda^2+x+y+\frac{(x-y)^2}{4\lambda^2}>0$, respectively. The following theorem establish the second-order distributional expansions of normalized maxima with $\lambda \in (0,\infty)$ and $\lambda^2+x+y+\frac{(x-y)^2}{4\lambda^2}<0$.

Theorem 2.1. *Let $\{(\xi_{ni},\eta_{ni}), 1\le i\le n, n\ge 1\}$ be a triangular array satisfying (1.1) with $\rho_n \in (0,1)$. Assume that F has upper endpoint equal 1 and further $1-F(1-t^{-1}) \in 2RV_{-\alpha,\tau}$ with $t>0$, $\alpha>0$, $\tau\le 0$ and auxiliary function $A(t)$. If there exists c_n satisfying $\lim_{n\to\infty} c_n = 0$ such that (1.4) with $\lambda_n = \left(\frac{1-\rho_n}{2a_n}\right)^{\frac{1}{2}}$ and $\lambda \in (0,\infty)$. With fixed $x<0$ and $y<0$ satisfying $\lambda^2+x+y+\frac{(x-y)^2}{4\lambda^2}<0$, we have*

$$\begin{aligned}
&\mathbb{P}\left(M_{n1}\le 1+a_nx, M_{n2}\le 1+a_ny\right) - H_{\alpha+\frac{1}{2},\lambda}(x,y)\\
=\;& H_{\alpha+\frac{1}{2},\lambda}(x,y)\Bigg\{-\frac{1}{2}n^{-1}\left(|x|^{\alpha+\frac{1}{2}}\psi_\alpha\left(\frac{\lambda+\frac{y-x}{2\lambda}}{\sqrt{2|x|}}\right)+|y|^{\alpha+\frac{1}{2}}\psi_\alpha\left(\frac{\lambda+\frac{x-y}{2\lambda}}{\sqrt{2|y|}}\right)\right)^2\\
&+Q_{\alpha+\frac{1}{2},\lambda}(x,y)+o(a_n+c_n+A(a_n^{-1})+n^{-1})\Bigg\} \qquad (2.1)
\end{aligned}$$

for large n, where

$$\begin{aligned}
&Q_{\alpha+\frac{1}{2},\lambda}(x,y)\\
=\;& |x|^{\alpha+\frac{1}{2}}\Bigg\{-\frac{a_n|x|}{c_\alpha}\int_{-1}^{\frac{\lambda+\frac{y-x}{2\lambda}}{\sqrt{2|x|}}}(1-s^2)^{\alpha-1}s^2(\alpha+1-\frac{3}{2}\alpha s^2-s^2)ds\\
&-\frac{A(a_n^{-1})|x|^{-\tau}}{c_\alpha}\int_{-1}^{\frac{\lambda+\frac{y-x}{2\lambda}}{\sqrt{2|x|}}}(1-s^2)^{\alpha}\frac{(1-s^2)^{-\tau}-1}{\tau}ds+\psi_\alpha\left(\frac{\lambda+\frac{y-x}{2\lambda}}{\sqrt{2|x|}}\right)\Bigg(-A(a_n^{-1})\frac{|x|^{-\tau}-1}{\tau}\\
&+\frac{a_n}{c_\alpha}\int_{-1}^{1}(1-s^2)^{\alpha-1}s^2(\alpha+1-\frac{3}{2}\alpha s^2-s^2)ds+\frac{A(a_n^{-1})}{c_\alpha}\int_{-1}^{1}\frac{(1-s^2)^{\alpha-\tau}-(1-s^2)^{\alpha}}{\tau}ds\Bigg)
\end{aligned}$$

$$
-\frac{\left(1-\frac{\left(\lambda+\frac{y-x}{2\lambda}\right)^2}{2|x|}\right)^{\alpha}}{c_\alpha\sqrt{2|x|}}\left(c_n\gamma\left(1-\frac{y-x}{2\lambda^2}\right)-\frac{a_n}{2}\left(\lambda(y-x)+\frac{x(y-x)}{\lambda}+\frac{3(y-x)^2}{4\lambda}+\frac{(y-x)^3}{8\lambda^3}\right)\right)\Bigg\}
$$

$$
+|y|^{\alpha+\frac{1}{2}}\Bigg\{-\frac{a_n|y|}{c_\alpha}\int_{-1}^{\frac{\lambda+\frac{x-y}{2\lambda}}{\sqrt{2|y|}}}(1-s^2)^{\alpha-1}s^2(\alpha+1-\frac{3}{2}\alpha s^2-s^2)ds
$$

$$
-\frac{A(a_n^{-1})|y|^{-\tau}}{c_\alpha}\int_{-1}^{\frac{\lambda+\frac{x-y}{2\lambda}}{\sqrt{2|y|}}}(1-s^2)^{\alpha}\frac{(1-s^2)^{-\tau}-1}{\tau}ds+\psi_\alpha\left(\frac{\lambda+\frac{x-y}{2\lambda}}{\sqrt{2|y|}}\right)\Bigg(-A(a_n^{-1})\frac{|y|^{-\tau}-1}{\tau}
$$

$$
+\frac{a_n}{c_\alpha}\int_{-1}^{1}(1-s^2)^{\alpha-1}s^2(\alpha+1-\frac{3}{2}\alpha s^2-s^2)ds+\frac{A(a_n^{-1})}{c_\alpha}\int_{-1}^{1}\frac{(1-s^2)^{\alpha-\tau}-(1-s^2)^{\alpha}}{\tau}ds\Bigg)
$$

$$
-\frac{\left(1-\frac{\left(\lambda+\frac{x-y}{2\lambda}\right)^2}{2|y|}\right)^{\alpha}}{c_\alpha\sqrt{2|y|}}\left(c_n\gamma\left(1-\frac{x-y}{2\lambda^2}\right)+\frac{a_n}{2}\left(\lambda(y-x)+\frac{x(y-x)}{\lambda}+\frac{(y-x)^2}{4\lambda}+\frac{(y-x)^3}{8\lambda^3}\right)\right)\Bigg\}
$$

$$
+o\left(a_n+c_n+A(a_n^{-1})\right),
$$

where $c_\alpha=\int_0^1(1-s)^\alpha s^{-\frac{1}{2}}ds=\frac{\Gamma(\alpha+1)\sqrt{\pi}}{\Gamma(\alpha+\frac{3}{2})}$.

For the case of $\lambda\in(0,\infty)$ with $\lambda^2+x+y+\frac{(x-y)^2}{4\lambda^2}=0$, restrictions on x and y may be needed. There are three cases to be considered: i), $x<0$ with $y=-\left(\sqrt{2}\lambda-\sqrt{-x}\right)^2$, and $y<0$ with $x=-\left(\sqrt{2}\lambda-\sqrt{-y}\right)^2$; ii), $x<0$ with $y=-(\sqrt{2}\lambda+\sqrt{-x})^2$ and, iii), $y<0$ with $x=-(\sqrt{2}\lambda+\sqrt{-y})^2$. The asymptotics are different and we present the results in turn.

Theorem 2.2. *Let* $\{(\xi_{ni},\eta_{ni}),1\le i\le n,n\ge1\}$ *be a triangular array satisfying* (1.1) *with* $\rho_n\in(0,1)$. *Assume that* F *has upper endpoint equal 1 and further* $1-F(1-t^{-1})\in 2RV_{-\alpha,\tau}$ *with* $t>0$, $\alpha>0$, $\tau\le0$ *and auxiliary function* $A(t)$. *Assuming that* (1.4) *holds with* $\lambda\in(0,\infty)$. *For* $x<0$ *with* $y=-\left(\sqrt{2}\lambda-\sqrt{-x}\right)^2$, *and* $y<0$ *with* $x=-\left(\sqrt{2}\lambda-\sqrt{-y}\right)^2$, *we have*

(i) if $\lim_{n\to\infty}\frac{c_n}{a_n}=k\in\mathbb{R}$, *then*

$$
\begin{aligned}
&\mathbb{P}\left(M_{n1}\le1+a_nx,M_{n2}\le1+a_ny\right)-H_{\alpha+\frac{1}{2},\lambda}(x,y)\\
=\;& H_{\alpha+\frac{1}{2},\lambda}(x,y)\Bigg\{-|x|^{\alpha+\frac{1}{2}}\left(\frac{a_n(|x|-1)}{c_\alpha}\int_0^1(1-s)^{\alpha-1}s^{\frac{1}{2}}\left(\alpha+1-\frac{3}{2}\alpha s-s\right)ds\right.\\
&\left.+A(a_n^{-1})\left(\frac{|x|^{-\tau}-1}{\tau}+\frac{|x|^{-\tau}-1}{c_\alpha}\int_0^1\frac{(1-s)^{\alpha-\tau}-(1-s)^{\alpha}}{\tau}s^{-\frac{1}{2}}ds\right)\right)\\
&-|y|^{\alpha+\frac{1}{2}}\left(\frac{a_n(|y|-1)}{c_\alpha}\int_0^1(1-s)^{\alpha-1}s^{\frac{1}{2}}\left(\alpha+1-\frac{3}{2}\alpha s-s\right)ds\right.
\end{aligned}
$$

$$+A(a_n^{-1})\left(\frac{|y|^{-\tau}-1}{\tau}+\frac{|y|^{-\tau}-1}{c_\alpha}\int_0^1\frac{(1-s)^{\alpha-\tau}-(1-s)^{\alpha}}{\tau}s^{-\frac{1}{2}}ds\right)\Bigg)$$
$$-\frac{1}{2}n^{-1}\left(|x|^{\alpha+\frac{1}{2}}+|y|^{\alpha+\frac{1}{2}}\right)^2+o\left(a_n+n^{-1}+A(a_n^{-1})\right)\Bigg\} \tag{2.2}$$

for large n;

(ii) if $a_n = o(c_n)$, then

$$\begin{aligned}
&\mathbb{P}\left(M_{n1}\le 1+a_nx, M_{n2}\le 1+a_ny\right)-H_{\alpha+\frac{1}{2},\lambda}(x,y)\\
=\;& H_{\alpha+\frac{1}{2},\lambda}(x,y)\Bigg\{-|x|^{\alpha+\frac{1}{2}}A(a_n^{-1})\left(\frac{|x|^{-\tau}-1}{\tau}+\frac{|x|^{-\tau}-1}{c_\alpha}\int_0^1\frac{(1-s)^{\alpha-\tau}-(1-s)^{\alpha}}{\tau}s^{-\frac{1}{2}}ds\right)\\
&-|y|^{\alpha+\frac{1}{2}}A(a_n^{-1})\left(\frac{|y|^{-\tau}-1}{\tau}+\frac{|y|^{-\tau}-1}{c_\alpha}\int_0^1\frac{(1-s)^{\alpha-\tau}-(1-s)^{\alpha}}{\tau}s^{-\frac{1}{2}}ds\right)\\
&-\frac{1}{2}n^{-1}\left(|x|^{\alpha+\frac{1}{2}}+|y|^{\alpha+\frac{1}{2}}\right)^2+o\left(c_n+n^{-1}+A(a_n^{-1})\right)\Bigg\}
\end{aligned} \tag{2.3}$$

for large n.

Theorem 2.3. *Let $\{(\xi_{ni},\eta_{ni}), 1\le i\le n, n\ge 1\}$ be a triangular array satisfying (1.1) with $\rho_n\in(0,1)$. Assume that F has upper endpoint equal 1 and further $1-F(1-t^{-1})\in 2RV_{-\alpha,\tau}$ with $t>0$ $\alpha>0$, $\tau\le 0$ and auxiliary function $A(t)$. Assuming that (1.4) holds with $\lambda\in(0,\infty)$. For $x<0$ with $y=-(\sqrt{2}\lambda+\sqrt{-x})^2$, we have*

(i) if $\lim_{n\to\infty}\frac{c_n}{a_n}=k\in\mathbb{R}$, then

$$\begin{aligned}
&\mathbb{P}\left(M_{n1}\le 1+a_nx, M_{n2}\le 1+a_ny\right)-H_{\alpha+\frac{1}{2},\lambda}(x,y)\\
=\;& H_{\alpha+\frac{1}{2},\lambda}(x,y)\Bigg\{-|y|^{\alpha+\frac{1}{2}}\left(\frac{a_n(|y|-1)}{c_\alpha}\int_0^1(1-s)^{\alpha-1}s^{\frac{1}{2}}\left(\alpha+1-\frac{3}{2}\alpha s-s\right)ds\right.\\
&\left.+A(a_n^{-1})\left(\frac{|y|^{-\tau}-1}{\tau}+\frac{|y|^{-\tau}-1}{c_\alpha}\int_0^1\frac{(1-s)^{\alpha-\tau}-(1-s)^{\alpha}}{\tau}s^{-\frac{1}{2}}ds\right)\right)\\
&-\frac{1}{2}n^{-1}|y|^{2\alpha+1}+o\left(a_n+n^{-1}+A(a_n^{-1})\right)\Bigg\}
\end{aligned} \tag{2.4}$$

for large n;

(ii) if $a_n = o(c_n)$, then

$$\mathbb{P}\left(M_{n1}\le 1+a_nx, M_{n2}\le 1+a_ny\right)-H_{\alpha+\frac{1}{2},\lambda}(x,y)$$

$$= H_{\alpha+\frac{1}{2},\lambda}(x,y)\left\{-|y|^{\alpha+\frac{1}{2}}A(a_n^{-1})\left(\frac{|y|^{-\tau}-1}{\tau}+\frac{|y|^{-\tau}-1}{c_\alpha}\int_0^1\frac{(1-s)^{\alpha-\tau}-(1-s)^{\alpha}}{\tau}s^{-\frac{1}{2}}ds\right)\right.$$
$$\left.-\frac{1}{2}n^{-1}|y|^{2\alpha+1}+o\left(c_n+n^{-1}+A(a_n^{-1})\right)\right\} \tag{2.5}$$

for large n.

Theorem 2.4. *Let $\{(\xi_{ni},\eta_{ni}), 1\le i\le n, n\ge 1\}$ be a triangular array satisfying (1.1) with $\rho_n\in(0,1)$. Assume that F has upper endpoint equal 1 and further $1-F(1-t^{-1})\in 2RV_{-\alpha,\tau}$ with $t>0$ $\alpha>0$, $\tau\le 0$ and auxiliary function $A(t)$. Assuming that (1.4) holds with $\lambda\in(0,\infty)$. For $y<0$ with $x=-(\sqrt{2}\lambda+\sqrt{-y})^2$, we have*

(i) if $\lim_{n\to\infty}\frac{c_n}{a_n}=k\in\mathbb{R}$, then

$$\begin{aligned}&\mathbb{P}\left(M_{n1}\le 1+a_nx, M_{n2}\le 1+a_ny\right)-H_{\alpha+\frac{1}{2},\lambda}(x,y)\\ =\ & H_{\alpha+\frac{1}{2},\lambda}(x,y)\left\{-|x|^{\alpha+\frac{1}{2}}\left(\frac{a_n(|x|-1)}{c_\alpha}\int_0^1(1-s)^{\alpha-1}s^{\frac{1}{2}}\left(\alpha+1-\frac{3}{2}\alpha s-s\right)ds\right.\right.\\ &\left.\left.+A(a_n^{-1})\left(\frac{|x|^{-\tau}-1}{\tau}+\frac{|x|^{-\tau}-1}{c_\alpha}\int_0^1\frac{(1-s)^{\alpha-\tau}-(1-s)^{\alpha}}{\tau}s^{-\frac{1}{2}}ds\right)\right)\right.\\ &\left.-\frac{1}{2}n^{-1}|x|^{2\alpha+1}+o\left(a_n+n^{-1}+A(a_n^{-1})\right)\right\}\end{aligned} \tag{2.6}$$

for large n;

(ii) if $a_n=o(c_n)$, then

$$\begin{aligned}&\mathbb{P}\left(M_{n1}\le 1+a_nx, M_{n2}\le 1+a_ny\right)-H_{\alpha+\frac{1}{2},\lambda}(x,y)\\ =\ & H_{\alpha+\frac{1}{2},\lambda}(x,y)\left\{-|x|^{\alpha+\frac{1}{2}}A(a_n^{-1})\left(\frac{|x|^{-\tau}-1}{\tau}+\frac{|x|^{-\tau}-1}{c_\alpha}\int_0^1\frac{(1-s)^{\alpha-\tau}-(1-s)^{\alpha}}{\tau}s^{-\frac{1}{2}}ds\right)\right.\\ &\left.-\frac{1}{2}n^{-1}|x|^{2\alpha+1}+o\left(c_n+n^{-1}+A(a_n^{-1})\right)\right\}\end{aligned} \tag{2.7}$$

for large n.

For $x<0, y<0$, note that $\lambda^2+x+y+\frac{(x-y)^2}{4\lambda^2}>0$ ensures that $\frac{\left(\lambda+\frac{y-x}{2\lambda}\right)^2}{-2x}>1$ and $\frac{\left(\lambda+\frac{x-y}{2\lambda}\right)^2}{-2y}>1$ all hold. Since $\lambda+\frac{y-x}{2\lambda}<-\sqrt{-2x}$ and $\lambda+\frac{x-y}{2\lambda}<-\sqrt{-2y}$ have no intersection, we can deal with the case of $\lambda\in(0,\infty)$ and $\lambda^2+x+y+\frac{(x-y)^2}{4\lambda^2}>0$ by the following three cases: (i) $\lambda+\frac{y-x}{2\lambda}>\sqrt{-2x}, \lambda+\frac{x-y}{2\lambda}>\sqrt{-2y}$; (ii) $\lambda+\frac{y-x}{2\lambda}<-\sqrt{-2x}, \lambda+\frac{x-y}{2\lambda}>\sqrt{-2y}$; (iii) $\lambda+\frac{y-x}{2\lambda}>\sqrt{-2x}, \lambda+\frac{x-y}{2\lambda}<-\sqrt{-2y}$.

Theorem 2.5. *Let* $\{(\xi_{ni}, \eta_{ni}), 1 \le i \le n, n \ge 1\}$ *be a triangular array satisfying* (1.1) *with* $\rho_n \in (0,1)$. *Assume that* F *has upper endpoint equal 1 and further* $1 - F(1 - t^{-1}) \in 2RV_{-\alpha,\tau}$ *with* $t > 0$, $\alpha > 0$, $\tau \le 0$ *and auxiliary function* $A(t)$. *If* (1.2) *holds with* $\lambda \in (0, \infty)$, *then*

(i) *if* $\lambda + \frac{y-x}{2\lambda} > \sqrt{-2x}, \lambda + \frac{x-y}{2\lambda} > \sqrt{-2y}$, *then* (2.2) *holds;*

(ii) *if* $\lambda + \frac{y-x}{2\lambda} < -\sqrt{-2x}, \lambda + \frac{x-y}{2\lambda} > \sqrt{-2y}$, *then* (2.4) *holds;*

(iii) *if* $\lambda + \frac{y-x}{2\lambda} > \sqrt{-2x}, \lambda + \frac{x-y}{2\lambda} < -\sqrt{-2y}$, *then* (2.6) *holds.*

To end this section, under some other additional condition related to ρ_n, we can investigate two extreme cases: $\lambda = 0$ and $\lambda = \infty$. For the case of $\lambda = 0$, we have the following results.

Theorem 2.6. *Let* $\{(\xi_{ni}, \eta_{ni}), 1 \le i \le n, n \ge 1\}$ *be a triangular array satisfying* (1.1) *with* $\rho_n \in (0,1]$. *Assume that* F *has upper endpoint equal 1 and further* $1 - F(1 - t^{-1}) \in 2RV_{-\alpha,\tau}$ *with* $t > 0$, $\alpha > 0$, $\tau \le 0$ *and auxiliary function* $A(t)$. *Then,*

(i), *if* $\rho_n = 1$, *then*

$$\begin{aligned}
& \mathbb{P}\left(M_{n1} \le 1 + a_n x, M_{n2} \le 1 + a_n y\right) - H_{\alpha+\frac{1}{2},0}(x,y) \\
= \; & H_{\alpha+\frac{1}{2},0}(x,y) \Bigg\{ -|\min(x,y)|^{\alpha+\frac{1}{2}} \left(\frac{a_n(|\min(x,y)| - 1)}{c_\alpha} \int_0^1 (1-s)^{\alpha-1} s^{\frac{1}{2}} \left(\alpha + 1 - \frac{3}{2}\alpha s - s \right) ds \right. \\
& \left. + A(a_n^{-1}) \left(\frac{|\min(x,y)|^{-\tau} - 1}{\tau} + \frac{|\min(x,y)|^{-\tau} - 1}{c_\alpha} \int_0^1 \frac{(1-s)^{\alpha-\tau} - (1-s)^\alpha}{\tau} s^{-\frac{1}{2}} ds \right) \right) \\
& - \frac{1}{2} n^{-1} |\min(x,y)|^{2\alpha+1} + o\left(a_n + n^{-1} + A(a_n^{-1})\right) \Bigg\} \qquad (2.8)
\end{aligned}$$

for large n;

(ii), *if* $\rho_n \in (0,1)$, *further assume that* $\lim_{n\to\infty} \frac{1-\rho_n}{a_n} = 0$ *and* $\lim_{n\to\infty} \frac{a_n^{5/3}}{1-\rho_n} = 0$, *then* (2.8) *also holds.*

For the case of $\lambda = \infty$, we have the following results.

Theorem 2.7. *Let* $\{(\xi_{ni}, \eta_{ni}), 1 \le i \le n, n \ge 1\}$ *be a triangular array satisfying* (1.1) *with* $\rho_n \in [-1,1)$. *Assume that* F *has upper endpoint equal 1 and further* $1 - F(1 - t^{-1}) \in 2RV_{-\alpha,\tau}$ *with* $t > 0$, $\alpha > 0$, $\tau \le 0$. *Then for* $\lambda = \infty$,

(i) *if* $\rho_n \to 1$, *further assume that* $\lim_{n\to\infty} \frac{1-\rho_n}{a_n} = \infty$ *and* $\lim_{n\to\infty} \frac{(1-\rho_n)^3}{a_n} = 0$, *then* (2.2) *holds;*

(ii) *if* $\rho_n \to d \in [-1, 1)$, *then* (2.2) *also holds.*

Example 2.1. *Consider a triangular array* $\{(\xi_{ni}, \eta_{ni}), 1 \le i \le n, n \ge 1\}$ *satisfying* (1.1) *with almost surely positive random radius* R *being Beta distributed with parameters* $a, b > 0$. *One can check that*

$$1 - F(1 - x^{-1}) = \bar{F}(1 - x^{-1}) = \frac{1}{bB(a,b)} x^{-b} \left(1 - \frac{b(a-1)x^{-1}}{b+1}(1 + o(1))\right), \ x \to \infty.$$

Hence, $\bar{F} \in 2RV_{-b,-1}$ *with auxiliary function* $A(t) = \frac{b(a-1)}{b+1} t^{-1}$. *Note that* $a_n = 1 - G^{\leftarrow}(1 - n^{-1})$. *By using Lemma 3.2, we have*

$$\begin{aligned} n^{-1} &= 1 - G(1 - a_n) \\ &= \frac{a_n^{b+\frac{1}{2}} c_b}{\sqrt{2}\pi b B(a,b)} \left(1 + \left(\frac{1}{c_b}\int_0^1 (1-s)^{b-1} s^{\frac{1}{2}}(b + 1 - \frac{3}{2}bs - s)ds - \frac{b(a-1)}{b+1} \right.\right. \\ &\quad \left.\left. + \frac{b(a-1)}{(b+1)c_b}\int_0^1 (1-s)^b s^{\frac{1}{2}} ds\right) a_n + o(a_n)\right) \end{aligned}$$

for large n, *where* $c_b = \int_0^1 (1-s)^b s^{-\frac{1}{2}} ds$. *So we can get*

$$a_n = \left(\frac{bB(a,b)\sqrt{2}\pi}{nc_b}\right)^{\frac{1}{b+\frac{1}{2}}} - \frac{m}{b+\frac{1}{2}} \left(\frac{bB(a,b)\sqrt{2}\pi}{nc_b}\right)^{\frac{2}{b+\frac{1}{2}}} (1 + o(1))$$

with

$$m = \frac{1}{c_b}\int_0^1 (1-s)^{b-1} s^{\frac{1}{2}}(b + 1 - \frac{3}{2}s - s)ds - \frac{b(a-1)}{b+1} + \frac{b(a-1)}{(b+1)c_b}\int_0^1 (1-s)^b s^{\frac{1}{2}} ds.$$

Assume that $\rho_n = 1 - n^{-\frac{1}{b+\frac{1}{2}}}$, *then the second-order condition* (1.4) *holds with*

$$\lambda = \frac{1}{\sqrt{2}} \left(\frac{bB(a,b)\sqrt{2}\pi}{c_b}\right)^{-\frac{1}{2b+1}}, \quad c_n = n^{-\frac{1}{b+\frac{1}{2}}}, \quad \text{and} \quad \gamma = \frac{m}{2\sqrt{2}(b+\frac{1}{2})} \left(\frac{bB(a,b)\sqrt{2}\pi}{c_b}\right)^{\frac{1}{2b+1}}.$$

For $x<0, y<0$ with $\lambda^2+x+y+\frac{(x-y)^2}{4\lambda^2}\leq 0$, from Theorem 2.1-2.4, it follows that

$$
\begin{aligned}
&\mathbb{P}\left(M_{n1}\leq 1+a_n x, M_{n2}\leq 1+a_n y\right)-H_{b,\lambda}(x,y)\\
=\;& H_{b,\lambda}(x,y)\left\{\left(\frac{bB(a,b)\sqrt{2}\pi}{nc_b}\right)^{\frac{1}{b+\frac{1}{2}}}\left[|x|^{b+\frac{1}{2}}\left(-\frac{|x|}{c_b}\int_{-1}^{\frac{\lambda+\frac{y-x}{2\lambda}}{\sqrt{2|x|}}}(1-s^2)^{b-1}s^2\left(b+1-\frac{3}{2}bs^2-s^2\right)ds\right.\right.\right.\\
&+\frac{b(a-1)}{c_b(b+1)}|x|\int_{-1}^{\frac{\lambda+\frac{y-x}{2\lambda}}{\sqrt{2|x|}}}\left((1-s^2)^{b+1}-(1-s^2)^b\right)ds+\psi_b\left(\frac{\lambda+\frac{y-x}{2\lambda}}{\sqrt{2|x|}}\right)\left(\frac{b(a-1)}{b+1}(|x|-1)\right.\\
&\left.+\frac{1}{c_b}\int_{-1}^{1}(1-s^2)^{b-1}s^2\left(b+1-\frac{3}{2}bs^2-s^2\right)ds-\frac{b(a-1)}{c_b(b+1)}\int_{-1}^{1}\left((1-s^2)^{b+1}-(1-s^2)^b\right)ds\right)\\
&-\frac{1}{c_b\sqrt{2|x|}}\left(1-\frac{\left(\lambda+\frac{y-x}{2\lambda}\right)^2}{2|x|}\right)^b\left(\gamma\left(1-\frac{y-x}{2\lambda^2}\right)\left(\frac{c_b}{bB(a,b)\sqrt{2}\pi}\right)^{\frac{1}{b+\frac{1}{2}}}\right.\\
&\left.\left.-\frac{1}{2}\left(\lambda(y-x)+\frac{(y-x)^3}{8\lambda^3}+\frac{(y-x)^2}{4\lambda}+\frac{x(y-x)}{\lambda}\right)\right)\right)\\
&+|y|^{b+\frac{1}{2}}\left(-\frac{|y|}{c_b}\int_{-1}^{\frac{\lambda+\frac{x-y}{2\lambda}}{\sqrt{2|y|}}}(1-s^2)^{b-1}s^2\left(b+1-\frac{3}{2}bs^2-s^2\right)ds\right.\\
&+\frac{b(a-1)}{c_b(b+1)}|y|\int_{-1}^{\frac{\lambda+\frac{x-y}{2\lambda}}{\sqrt{2|y|}}}\left((1-s^2)^{b+1}-(1-s^2)^b\right)ds+\psi_b\left(\frac{\lambda+\frac{x-y}{2\lambda}}{\sqrt{2|y|}}\right)\left(\frac{b(a-1)}{b+1}(|y|-1)\right.\\
&\left.+\frac{1}{c_b}\int_{-1}^{1}(1-s^2)^{b-1}s^2\left(b+1-\frac{3}{2}bs^2-s^2\right)ds-\frac{b(a-1)}{c_b(b+1)}\int_{-1}^{1}\left((1-s^2)^{b+1}-(1-s^2)^b\right)ds\right)\\
&-\frac{1}{c_b\sqrt{2|y|}}\left(1-\frac{\left(\lambda+\frac{x-y}{2\lambda}\right)^2}{2|y|}\right)^b\left(\gamma\left(1-\frac{x-y}{2\lambda^2}\right)\left(\frac{c_b}{bB(a,b)\sqrt{2}\pi}\right)^{\frac{1}{b+\frac{1}{2}}}\right.\\
&\left.\left.\left.+\frac{1}{2}\left(\lambda(y-x)+\frac{(y-x)^3}{8\lambda^3}+\frac{(y-x)^2}{4\lambda}+\frac{x(y-x)}{\lambda}\right)\right)\right)\right]\\
&-\frac{1}{2}n^{-1}\left(|x|^{b+\frac{1}{2}}\psi_b\left(\frac{\lambda+\frac{y-x}{2\lambda}}{\sqrt{2|x|}}\right)+|y|^{b+\frac{1}{2}}\psi_b\left(\frac{\lambda+\frac{x-y}{2\lambda}}{\sqrt{2|y|}}\right)\right)^2+o\left(n^{-1}+n^{-\frac{1}{b+\frac{1}{2}}}\right)\Bigg\}.
\end{aligned}
$$

For $x<0$, $y<0$ with $\lambda^2+x+y+\frac{(x-y)^2}{4\lambda^2}>0$, using Theorem 2.5 we can get

$$
\begin{aligned}
&\mathbb{P}\left(M_{n1}\leq 1+a_n x, M_{n2}\leq 1+a_n y\right)-H_{b,\lambda}(x,y)\\
=\;& H_{b,\lambda}(x,y)\left\{\left(\frac{bB(a,b)\sqrt{2}\pi}{nc_b}\right)^{\frac{1}{b+\frac{1}{2}}}\left[|x|^{b+\frac{1}{2}}\left(-\frac{|x|}{c_b}\int_{-1}^{sgn\left(\frac{\lambda+\frac{y-x}{2\lambda}}{\sqrt{2|x|}}\right)}(1-s^2)^{b-1}s^2\left(b+1-\frac{3}{2}bs^2-s^2\right)ds\right.\right.\right.\\
&+\frac{b(a-1)}{c_b(b+1)}|x|\int_{-1}^{sgn\left(\frac{\lambda+\frac{y-x}{2\lambda}}{\sqrt{2|x|}}\right)}\left((1-s^2)^{b+1}-(1-s^2)^b\right)ds+\psi_b\left(\frac{\lambda+\frac{y-x}{2\lambda}}{\sqrt{2|x|}}\right)\left(\frac{b(a-1)}{b+1}(|x|-1)\right.
\end{aligned}
$$

$$+\frac{1}{c_b}\int_{-1}^{1}(1-s^2)^{b-1}s^2\left(b+1-\frac{3}{2}bs^2-s^2\right)ds-\frac{b(a-1)}{c_b(b+1)}\int_{-1}^{1}\left((1-s^2)^{b+1}-(1-s^2)^b\right)ds\Bigg)\Bigg)$$
$$+|y|^{b+\frac{1}{2}}\left(-\frac{|y|}{c_b}\int_{-1}^{sgn\left(\frac{\lambda+\frac{x-y}{2\lambda}}{\sqrt{2|y|}}\right)}(1-s^2)^{b-1}s^2\left(b+1-\frac{3}{2}bs^2-s^2\right)ds\right.$$
$$+\frac{b(a-1)}{c_b(b+1)}|y|\int_{-1}^{sgn\left(\frac{\lambda+\frac{x-y}{2\lambda}}{\sqrt{2|y|}}\right)}\left((1-s^2)^{b+1}-(1-s^2)^b\right)ds+\psi_b\left(\frac{\lambda+\frac{x-y}{2\lambda}}{\sqrt{2|y|}}\right)\left(\frac{b(a-1)}{b+1}(|y|-1)\right.$$
$$\left.\left.+\frac{1}{c_b}\int_{-1}^{1}(1-s^2)^{b-1}s^2\left(b+1-\frac{3}{2}bs^2-s^2\right)ds-\frac{b(a-1)}{c_b(b+1)}\int_{-1}^{1}\left((1-s^2)^{b+1}-(1-s^2)^b\right)ds\right)\right)\Bigg]$$
$$-\frac{1}{2}n^{-1}\left(|x|^{b+\frac{1}{2}}\psi_b\left(\frac{\lambda+\frac{y-x}{2\lambda}}{\sqrt{2|x|}}\right)+|y|^{b+\frac{1}{2}}\psi_b\left(\frac{\lambda+\frac{x-y}{2\lambda}}{\sqrt{2|y|}}\right)\right)^2+o\left(n^{-1}+n^{-\frac{1}{b+\frac{1}{2}}}\right)\Bigg\}$$

with the condition (1.2) *holding, where sgn(·) is sign function.*

3 Auxiliary lemmas

In order to prove our main results, we give some auxiliary lemmas in this section. For simplicity, in the sequel, let $t_n(x)=1+a_nx$, where $x<0$ and $a_n=1-G^{-1}(1-n^{-1})$. The first lemma is about Drees' type inequalities for the second-order regular varying functions $2RV_{\alpha,\tau}$, cf. de Haan and Ferreira (2006).

Lemma 3.1. *If $\chi\in 2RV_{\alpha,\tau}$ with auxiliary function $A_1(t),\alpha\in R$ and $\tau\le 0$, then for any $\varepsilon,\delta>0$, there exist an auxiliary function $A(t),A(t)\sim A_1(t)$ as $t\to\infty$, and $t_0=t_0(\varepsilon,\delta)>0$ such that for all $t,tx>t_0$,*

$$\left|\frac{\frac{\chi(tx)}{\chi(t)}-x^\alpha}{A(t)}-x^\alpha\frac{x^\tau-1}{\tau}\right|\le\varepsilon x^{\alpha+\tau}\max\left(x^\delta,x^{-\delta}\right).$$

Without loss of generality, we assume that auxiliary functions of $2RV$ functions are positive eventually in the following proofs. Before providing the distribution tail expansion of G, we need the following lemma.

Lemma 3.2. *Let Q be a Beta distributed random variable with positive parameters a and b. Assume that distribution function F with upper endpoint 1 satisfies $1-F(1-t^{-1})\in 2RV_{-\alpha,\tau}$ with $\alpha>$*

$0, \tau \leq 0$ *and auxiliary function* $A(t)$*. Then for large* n *we have*

$$\begin{aligned} & \mathbf{E}\left(1-F\left(t_n(x)(1-Q)^{-\frac{1}{2}}\right)\right) \\ = \quad & \frac{(2a_n|x|)^a\Gamma(a+b)}{\Gamma(a)\Gamma(b)}[1-F(t_n(x))]\left(\int_0^1(1-s)^\alpha s^{a-1}ds + A(a_n^{-1})|x|^{-\tau}\int_0^1(1-s)^\alpha\frac{(1-s)^{-\tau}-1}{\tau}s^{a-1}ds\right. \\ & \left.+a_n|x|\int_0^1(1-s)^{\alpha-1}s^a\left(\alpha-2(b-1)-\frac{3}{2}\alpha s+2(b-1)s\right)ds+o\left(a_n+A(a_n^{-1})\right)\right) \end{aligned} \tag{3.1}$$

with $x<0$*. Here,* $\frac{(1-y)^{-\tau}-1}{\tau}$ *is interpreted as* $-\log(1-y)$ *when* $\tau=0$*.*

Proof. In order to get the desired result, we first give the following inequalities for $0<x<\frac{1}{2}$:

$$1-\alpha x<(1-x)^\alpha\leq 1-\alpha x+\frac{\alpha(\alpha-1)}{2}x^2,\quad \alpha\geq 2, \tag{3.2}$$

$$1-\alpha x\leq(1-x)^\alpha\leq 1-\alpha x+2^{1-\alpha}\alpha(\alpha-1)x^2,\quad 1\leq\alpha<2, \tag{3.3}$$

$$1-\alpha x+2^{1-\alpha}\alpha(\alpha-1)x^2<(1-x)^\alpha<1-\alpha x,\quad 0<\alpha<1 \tag{3.4}$$

and

$$1+\alpha x<(1+x)^\alpha\leq 1+\alpha x+\left(\frac{3}{2}\right)^{\alpha-2}\frac{\alpha(\alpha-1)}{2}x^2,\quad \alpha\geq 2, \tag{3.5}$$

$$1+\alpha x\leq(1+x)^\alpha\leq 1+\alpha x+\frac{\alpha(\alpha-1)}{2}x^2,\quad 1\leq\alpha<2, \tag{3.6}$$

$$1+\alpha x+\frac{\alpha(\alpha-1)}{2}x^2<(1+x)^\alpha<1+\alpha x,\quad 0<\alpha<1. \tag{3.7}$$

Note that for $\beta>-1, 1<c<\alpha+1$ and large n, we have

$$\int_{1-\frac{(a_n|x|)^{\frac{1}{c}}}{2}}^{1-\frac{a_n|x|}{2}}\left(1-t_n(x)s-\frac{3}{2}a_n|x|t_n(x)s^2\right)^\alpha s^\beta ds$$

$$
\begin{aligned}
&= \int_{1-\frac{(a_n|x|)^{\frac{1}{c}}}{2}}^{1-\frac{a_n|x|}{2}} (1-s)^{\alpha} s^{\beta} \left(1+\frac{a_n|x|s\left(1-\frac{3}{2}s\right)+\frac{3}{2}a_n^2|x|^2s^2}{1-s}\right)^{\alpha} ds \\
&< \int_{1-\frac{(a_n|x|)^{\frac{1}{c}}}{2}}^{1-\frac{a_n|x|}{2}} (1-s)^{\alpha} s^{\beta} ds \\
&< \frac{(a_n|x|)^{\frac{\alpha}{c}}}{2^{\alpha}} \int_{1-\frac{(a_n|x|)^{\frac{1}{c}}}{2}}^{1-\frac{a_n|x|}{2}} s^{\beta} ds \\
&= \frac{(a_n|x|)^{\frac{\alpha+1}{c}}}{2^{\alpha+1}}(1+o(1)) = o(a_n),
\end{aligned} \tag{3.8}
$$

$$
\int_{1-\frac{(a_n|x|)^{\frac{1}{c}}}{2}}^{1} (1-s)^{\alpha-1} s^{\beta+1} ds < \int_{1-\frac{(a_n|x|)^{\frac{1}{c}}}{2}}^{1} (1-s)^{\alpha-1} ds = \frac{(a_n|x|)^{\frac{\alpha}{c}}}{\alpha 2^{\alpha}} = O(a_n^{\alpha/c}), \tag{3.9}
$$

$$
\int_{1-\frac{(a_n|x|)^{\frac{1}{c}}}{2}}^{1} (1-s)^{\alpha} s^{\beta} ds \le \frac{(a_n|x|)^{\frac{\alpha}{c}}}{2^{\alpha}(\beta+1)} \left(1-\left(1-\frac{(a_n|x|)^{\frac{1}{c}}}{2}\right)^{\beta+1}\right) = o(a_n) \tag{3.10}
$$

and

$$
\begin{aligned}
& a_n^2 \int_{\frac{2}{3}}^{1-\frac{(a_n|x|)^{\frac{1}{c}}}{2}} (1-s)^{\alpha-2} s^{\beta+2} \left(1-\frac{3}{2}s+\frac{3}{2}a_n|x|s\right)^2 ds \\
< \; & 9a_n^2 \int_{\frac{2}{3}}^{1-\frac{(a_n|x|)^{\frac{1}{c}}}{2}} (1-s)^{\alpha-2} ds \\
= \; & \frac{9a_n^2}{1-\alpha}\left(\frac{(a_n|x|)^{\frac{\alpha-1}{c}}}{2^{\alpha-1}} - 3^{1-\alpha}\right) \\
< \; & \frac{9a_n^{2+\frac{\alpha-1}{c}}|x|^{\frac{\alpha-1}{c}}}{(1-\alpha)2^{\alpha-1}} = o(a_n)
\end{aligned} \tag{3.11}
$$

for large n, if $0<\alpha<1$. From (3.4) and (3.9)-(3.11), it follows that

$$
\begin{aligned}
& \int_{\frac{2}{3}}^{1-\frac{(a_n|x|)^{\frac{1}{c}}}{2}} \left(1-t_n(x)s-\frac{3}{2}a_n|x|t_n(x)s^2\right)^{\alpha} s^{\beta} ds \\
< \; & \int_{\frac{2}{3}}^{1-\frac{(a_n|x|)^{\frac{1}{c}}}{2}} (1-s)^{\alpha} s^{\beta} ds + \alpha \int_{\frac{2}{3}}^{1-\frac{(a_n|x|)^{\frac{1}{c}}}{2}} (1-s)^{\alpha-1} s^{\beta+1} a_n|x| \left(1-\frac{3}{2}s+\frac{3}{2}a_n|x|s\right) ds \\
= \; & \int_{\frac{2}{3}}^{1} (1-s)^{\alpha} s^{\beta} ds + \alpha a_n|x| \int_{\frac{2}{3}}^{1} (1-s)^{\alpha-1} s^{\beta+1} \left(1-\frac{3}{2}s\right) ds + o(a_n)
\end{aligned} \tag{3.12}
$$

and

$$\begin{aligned}
&\int_{\frac{2}{3}}^{1-\frac{(a_n|x|)^{\frac{1}{c}}}{2}} \left(1 - t_n(x)s - \frac{3}{2}a_n|x|t_n(x)s^2\right)^{\alpha} s^{\beta} ds \\
> &\int_{\frac{2}{3}}^{1-\frac{(a_n|x|)^{\frac{1}{c}}}{2}} (1-s)^{\alpha} s^{\beta} ds + \alpha \int_{\frac{2}{3}}^{1-\frac{(a_n|x|)^{\frac{1}{c}}}{2}} (1-s)^{\alpha-1} s^{\beta+1} a_n|x| \left(1 - \frac{3}{2}s + \frac{3}{2}a_n|x|s\right) ds \\
&+2^{1-\alpha}\alpha(\alpha-1)a_n^2|x|^2 \int_{\frac{2}{3}}^{1-\frac{(a_n|x|)^{\frac{1}{c}}}{2}} (1-s)^{\alpha-2} s^{\beta+2} \left(1 - \frac{3}{2}s + \frac{3}{2}a_n|x|s\right)^2 ds \\
= &\int_{\frac{2}{3}}^{1} (1-s)^{\alpha} s^{\beta} ds + \alpha a_n|x| \int_{\frac{2}{3}}^{1} (1-s)^{\alpha-1} s^{\beta+1} \left(1 - \frac{3}{2}s\right) ds + o(a_n) \qquad (3.13)
\end{aligned}$$

for large n. Combining with (3.8), we can get

$$\begin{aligned}
&\int_{\frac{2}{3}}^{1-\frac{a_n|x|}{2}} \left(1 - t_n(x)s - \frac{3}{2}a_n|x|t_n(x)s^2\right)^{\alpha} s^{\beta} ds \\
= &\int_{\frac{2}{3}}^{1} (1-s)^{\alpha} s^{\beta} ds + \alpha a_n|x| \int_{\frac{2}{3}}^{1} (1-s)^{\alpha-1} s^{\beta+1} \left(1 - \frac{3}{2}s\right) ds + o(a_n), \qquad (3.14)
\end{aligned}$$

if $0 < \alpha < 1$.

Now we consider the cases of $\alpha \geq 1$. Since for large n and $\beta > -1$

$$\int_{1-\frac{a_n|x|}{2}}^{1} (1-s)^{\alpha-2} s^{\beta+2} ds \leq \int_{1-\frac{a_n|x|}{2}}^{1} (1-s)^{\alpha-2} ds = \frac{(a_n|x|)^{\alpha-1}}{(\alpha-1)2^{\alpha-1}} = O(a_n^{\alpha-1})$$

and

$$\int_{1-\frac{a_n|x|}{2}}^{1} (1-s)^{\alpha-1} s^{\beta+1} ds \leq \int_{1-\frac{a_n|x|}{2}}^{1} (1-s)^{\alpha-1} ds = \frac{(a_n|x|)^{\alpha}}{\alpha 2^{\alpha}} = o(a_n)$$

hold with $\alpha > 1$, and

$$\int_{1-\frac{a_n|x|}{2}}^{1} (1-s)^{\alpha} s^{\beta} ds \leq \left(\frac{a_n|x|}{2}\right)^{\alpha} \int_{1-\frac{a_n|x|}{2}}^{1} s^{\beta} ds = \frac{(a_n|x|)^{\alpha+1}}{2^{\alpha+1}}(1 + o(1)) = o(a_n)$$

holds with $\alpha > 0$, (3.14) also holds for the cases $\alpha \geq 1$ by using arguments similar to that of (3.12) and (3.13).

Similarly, by using (3.5)-(3.7) we have

$$\begin{aligned}
&\int_0^{\frac{2}{3}} \left(1 - t_n(x)s - \frac{3}{2}a_n|x|t_n(x)s^2\right)^{\alpha} s^{\beta} ds \\
= &\int_0^{\frac{2}{3}} (1-s)^{\alpha} s^{\beta} ds + \alpha a_n|x| \int_0^{\frac{2}{3}} (1-s)^{\alpha-1} s^{\beta+1} \left(1 - \frac{3}{2}s\right) ds + o(a_n)
\end{aligned}$$

for large n, which implies that

$$\begin{aligned}
&\int_0^{1-\frac{a_n|x|}{2}} \left(1 - t_n(x)s - \frac{3}{2}a_n|x|t_n(x)s^2\right)^{\alpha} s^{\beta} ds \\
= &\int_0^{1} (1-s)^{\alpha} s^{\beta} ds + \alpha a_n|x| \int_0^{1} (1-s)^{\alpha-1} s^{\beta+1} \left(1 - \frac{3}{2}s\right) ds + o(a_n) \qquad (3.15)
\end{aligned}$$

by combining with (3.14).

From Lemma 3.1, (3.15) and the following inequalities

$$1 + \frac{1}{2}s + \frac{3}{8}s^2 < (1-s)^{-\frac{1}{2}} < 1 + \frac{1}{2}s + \frac{3}{8}s^2 + \frac{5\sqrt{2}}{2}s^3, 0 < s < \frac{1}{2}$$

and

$$1 - (b-1)s < (1-s)^{b-1} < 1 - (b-1)s + 2^{2-b}(b-1)(b-2)s^2, 0 < s < \frac{1}{2}, b < 1,$$

we can get

$$\begin{aligned}
&\mathbf{E}\left(1 - F(t_n(x)(1-Q)^{-\frac{1}{2}})\right) \\
= &\frac{\Gamma(a+b)}{\Gamma(a)\Gamma(b)} \int_0^{2a_n|x|-a_n^2x^2} \left(1 - F\left(t_n(x)(1-s)^{-\frac{1}{2}}\right)\right) s^{a-1}(1-s)^{b-1} ds \\
\leq &\frac{\Gamma(a+b)}{\Gamma(a)\Gamma(b)} \int_0^{2a_n|x|-a_n^2x^2} \left(1 - F\left(t_n(x)\left(1 + \frac{1}{2}s + \frac{3}{8}s^2\right)\right)\right) \\
&\times s^{a-1}\left(1 - (b-1)s + 2^{2-b}(b-1)(b-2)s^2\right) ds \\
= &\frac{\Gamma(a+b)}{\Gamma(a)\Gamma(b)} (2a_n|x|)^a \int_0^{1-\frac{a_n|x|}{2}} \left(1 - F\left(t_n(x)\left(1 + a_n|x|s + \frac{3}{2}a_n^2x^2s^2\right)\right)\right) \\
&\times s^{a-1}\left(1 - 2a_n|x|(b-1)s + 2^{4-b}(b-1)(b-2)a_n^2x^2s^2\right) ds \\
\leq &\frac{\Gamma(a+b)}{\Gamma(a)\Gamma(b)} (2a_n|x|)^a \left(1 - F(t_n(x))\right) \\
&\times \left(\int_0^{1-\frac{a_n|x|}{2}} \left(1 - t_n(x)s - \frac{3}{2}a_n|x|t_n(x)s^2\right)^{\alpha} s^{a-1}\left(1 - 2a_n|x|(b-1)s + 2^{4-b}(b-1)(b-2)a_n^2x^2s^2\right) ds\right.
\end{aligned}$$

$$
\begin{aligned}
&+A(a_n^{-1}|x|^{-1})\int_0^{1-\frac{a_n|x|}{2}}\left(\left(1-t_n(x)s-\frac{3}{2}a_n|x|t_n(x)s^2\right)^{\alpha}\frac{\left(1-t_n(x)s-\frac{3}{2}a_n|x|t_n(x)s^2\right)^{-\tau}-1}{\tau}\right.\\
&\left.+\varepsilon\left(1-t_n(x)s-\frac{3}{2}a_n|x|t_n(x)s^2\right)^{\alpha-\tau-\varepsilon}\right)\times s^{a-1}\left(1-2a_n|x|(b-1)s+2^{4-b}(b-1)(b-2)a_n^2x^2s^2\right)ds\Bigg)\\
=\;&\frac{\Gamma(a+b)}{\Gamma(a)\Gamma(b)}(2a_n|x|)^a\left(1-F(t_n(x))\right)\left(\int_0^1(1-s)^{\alpha}s^{a-1}ds+A(a_n^{-1})|x|^{-\tau}\int_0^1(1-s)^{\alpha}\frac{(1-s)^{-\tau}-1}{\tau}s^{a-1}ds\right.\\
&\left.+a_n|x|\int_0^1(1-s)^{\alpha-1}s^a\left(\alpha-2(b-1)-\frac{3}{2}\alpha s+2(b-1)s\right)ds+o\left(a_n+A(a_n^{-1})\right)\right) \qquad (3.16)
\end{aligned}
$$

for large n. Similarly,

$$
\begin{aligned}
&\mathbf{E}\left(1-F\left(t_n(x)(1-Q)^{-\frac{1}{2}}\right)\right)\\
\geq\;&\frac{\Gamma(a+b)}{\Gamma(a)\Gamma(b)}\int_0^{2a_n|x|-a_n^2x^2}\left(1-F\left(t_n(x)\left(1+\frac{1}{2}s+\frac{3}{8}s^2+\frac{5\sqrt{2}}{2}s^3\right)\right)\right)s^{a-1}(1-(b-1)s)ds\\
\geq\;&\frac{\Gamma(a+b)}{\Gamma(a)\Gamma(b)}(2a_n|x|)^a\left(1-F(t_n(x))\right)\\
&\times\left(\int_0^{1-\frac{a_n|x|}{2}}\left(1-t_n(x)s-\frac{3}{2}a_n|x|t_n(x)s^2-20\sqrt{2}a_n^2x^2t_n(x)s^3\right)^{\alpha}s^{a-1}\left(1-2a_n|x|(b-1)s\right)ds\right.\\
&+A(a_n^{-1}|x|^{-1})\int_0^{1-\frac{a_n|x|}{2}}\left(\left(1-t_n(x)s-\frac{3}{2}a_n|x|t_n(x)s^2-20\sqrt{2}a_n^2x^2t_n(x)s^3\right)^{\alpha}\right.\\
&\times\frac{\left(1-t_n(x)s-\frac{3}{2}a_n|x|t_n(x)s^2-20\sqrt{2}a_n^2x^2t_n(x)s^3\right)^{-\tau}-1}{\tau}\\
&\left.\left.-\varepsilon\left(1-t_n(x)s-\frac{3}{2}a_n|x|t_n(x)s^2-20\sqrt{2}a_n^2x^2t_n(x)s^3\right)^{\alpha-\tau-\varepsilon}\right)\times s^{a-1}\left(1-2a_n|x|(b-1)s\right)ds\right)\\
=\;&\frac{\Gamma(a+b)}{\Gamma(a)\Gamma(b)}(2a_n|x|)^a\left(1-F(t_n(x))\right)\left(\int_0^1(1-s)^{\alpha}s^{a-1}ds+A(a_n^{-1})|x|^{-\tau}\int_0^1(1-s)^{\alpha}\frac{(1-s)^{-\tau}-1}{\tau}s^{a-1}ds\right.\\
&\left.+a_n|x|\int_0^1(1-s)^{\alpha-1}s^a\left(\alpha-2(b-1)-\frac{3}{2}\alpha s+2(b-1)s\right)ds+o\left(a_n+A(a_n^{-1})\right)\right). \qquad (3.17)
\end{aligned}
$$

Combining (3.16) and (3.17), we can derive (3.1), which complete the proof. □

Lemma 3.3. *Let (S_1, S_2) be a bivariate spherical random vector with almost surely positive random radius R with df F. Assume that F has upper endpoint 1 and further $1-F(1-t^{-1}) \in 2RV_{-\alpha,\tau}$ with $t>0, \alpha>0, \tau\leq 0$ and auxiliary function $A(t)$. Then $1-G(1-t^{-1}) \in 2RV_{-\alpha-\frac{1}{2},\max(-1,\tau)}$, where G is the df of S_1.*

Proof. By Corollary 12.1.1 of Berman (1992) and arguments similar to Lemma 3.2, we can get

$$
1-G(1-(t|x|)^{-1})
$$

$$
\begin{aligned}
&= \frac{1}{2}\mathbf{E}\left(1-F\left((1-(t|x|)^{-1})(1-Q)^{-\frac{1}{2}}\right)\right)\\
&= \frac{(2t^{-1}|x|^{-1})^{\frac{1}{2}}}{2\pi}\left(1-F\left(1-(t|x|)^{-1}\right)\right)\\
&\quad\times\left\{\int_0^1(1-s)^{\alpha}s^{-\frac{1}{2}}ds+(t|x|)^{-1}\int_0^1(1-s)^{\alpha-1}s^{\frac{1}{2}}\left(\alpha+1-\frac{3}{2}\alpha s-s\right)ds\right.\\
&\quad\left.+A(t)|x|^{\tau}\int_0^1(1-s)^{\alpha}\frac{(1-s)^{-\tau}-1}{\tau}s^{-\frac{1}{2}}ds+o(t^{-1}+A(t))\right\} \qquad (3.18)
\end{aligned}
$$

for large t and $x<0$. Here Q is a Beta distributed random variable with parameters $1/2, 1/2$. Hence,

$$
\begin{aligned}
\frac{1-G(1-(t|x|)^{-1})}{1-G(1-t^{-1})} &= |x|^{-\frac{1}{2}-\alpha}\left(1+\frac{t^{-1}(|x|^{-1}-1)}{c_\alpha}\int_0^1(1-s)^{\alpha-1}s^{\frac{1}{2}}\left(\alpha+1-\frac{3}{2}\alpha s-s\right)ds\right.\\
&\quad\left.+A(t)\left(\frac{|x|^{\tau}-1}{\tau}+\frac{|x|^{\tau}-1}{c_\alpha}\int_0^1(1-s)^{\alpha}\frac{(1-s)^{-\tau}-1}{\tau}s^{-\frac{1}{2}}ds\right)+o(t^{-1}+A(t))\right) \qquad (3.19)
\end{aligned}
$$

for large t, where $c_\alpha=\int_0^1(1-s)^{\alpha}s^{-\frac{1}{2}}ds$. So $1-G(1-t^{-1})\in 2RV_{-\alpha-\frac{1}{2},\max(-1,\tau)}$. □

4 Proofs of main results

The aim of this section is to prove our main results.

Proof of Theorem 2.1. Note that $\arccos z=\sqrt{2(1-z)}\left(1+\frac{1-z}{12}+o(1-z)\right)$ for $z\to 1$. By (1.4) and the Taylor expansion with the Lagrange remainder, we have

$$
\begin{aligned}
\frac{\frac{t_n(y)}{t_n(x)}-\rho_n}{\sqrt{a_n}\sin\tau_n} &= \frac{1-\rho_n+\frac{a_n(y-x)}{1+a_nx}}{\sqrt{a_n}\left(\sqrt{2(1-\rho_n)}\left(1+\frac{1}{12}(1-\rho_n)+o(1-\rho_n)\right)-\frac{(\sqrt{2(1-\rho_n)})^3}{6}(1+o(1))\right)}\\
&= \lambda+\frac{y-x}{2\lambda}+\gamma\left(1-\frac{y-x}{2\lambda^2}\right)c_n+\left(\frac{\lambda^3}{2}+\frac{(y-x)\lambda}{4}-\frac{x(y-x)}{2\lambda}\right)a_n+o(a_n+c_n) \qquad (4.1)
\end{aligned}
$$

for large n, where $\tau_n=\arccos\rho_n$. Then for $\beta_n(x,y)=\arctan\left(\frac{\frac{y}{x}-\rho_n}{\sin\tau_n}\right)$, we can get

$$
\begin{aligned}
&\frac{\sin\beta_n\left(t_n(x),t_n(y)\right)}{\sqrt{a_n}}\\
&= \frac{\beta_n\left(t_n(x),t_n(y)\right)}{\sqrt{a_n}}-\frac{\beta_n^3\left(t_n(x),t_n(y)\right)}{6\sqrt{a_n}}(1+o(1))\\
&= \lambda+\frac{y-x}{2\lambda}+\gamma\left(1-\frac{y-x}{2\lambda^2}\right)c_n
\end{aligned}
$$

$$-\frac{a_n}{2}\left(\lambda(y-x)+\frac{x(y-x)}{\lambda}+\frac{3(y-x)^2}{4\lambda}+\frac{(y-x)^3}{8\lambda^3}\right)+o(a_n+c_n) \tag{4.2}$$

for large n.

First we consider the case of $\lambda+\frac{y-x}{2\lambda}\geq 0$ with $\lambda^2+x+y+\frac{(x-y)^2}{4\lambda^2}<0$. Combining (4.2) and Taylor's Theorem, we can get

$$\begin{aligned}
&\int_{\frac{\sin^2\beta_n(t_n(x),t_n(y))}{2a_n|x|}}^{1-\frac{a_n|x|}{2}}\left(1-t_n(x)s-\frac{3}{2}a_n|x|t_n(x)s^2\right)^{\alpha}s^{\beta}ds\\
=\;&\int_{\frac{\left(\lambda+\frac{y-x}{2\lambda}\right)^2}{2|x|}}^{1-\frac{a_n|x|}{2}}\left(1-t_n(x)s-\frac{3}{2}a_n|x|t_n(x)s^2\right)^{\alpha}s^{\beta}ds-\left(\frac{\sin^2\beta_n(t_n(x),t_n(y))}{2a_n|x|}-\frac{\left(\lambda+\frac{y-x}{2\lambda}\right)^2}{2|x|}\right)\\
&\times\left(1-\frac{1+a_nx}{2|x|}\left(\lambda+\frac{y-x}{2\lambda}\right)^2-\frac{3}{2}a_n|x|\frac{\left(\lambda+\frac{y-x}{2\lambda}\right)^4}{4x^2}(1+a_nx)\right)^{\alpha}\frac{\left(\lambda+\frac{y-x}{2\lambda}\right)^{2\beta}}{(2|x|)^{\beta}}(1+o(1))\\
=\;&\int_{\frac{\left(\lambda+\frac{y-x}{2\lambda}\right)^2}{2|x|}}^{1}(1-s)^{\alpha}s^{\beta}ds+\alpha a_n|x|\int_{\frac{\left(\lambda+\frac{y-x}{2\lambda}\right)^2}{2|x|}}^{1}(1-s)^{\alpha-1}s^{\beta+1}\left(1-\frac{3}{2}s\right)ds\\
&-\frac{\left(\lambda+\frac{y-x}{2\lambda}\right)^{2\beta+1}}{2^{\beta}|x|^{\beta+1}}\left(1-\frac{\left(\lambda+\frac{y-x}{2\lambda}\right)^2}{2|x|}\right)^{\alpha}\left\{c_n\gamma\left(1-\frac{y-x}{2\lambda^2}\right)\right.\\
&\left.-\frac{a_n}{2}\left(\lambda(y-x)+\frac{x(y-x)}{\lambda}+\frac{3(y-x)^2}{4\lambda}+\frac{(y-x)^3}{8\lambda^3}\right)\right\}+o(a_n+c_n)
\end{aligned}$$

for large n and $\beta>-1$, since

$$\begin{aligned}
&\int_{\frac{\left(\lambda+\frac{y-x}{2\lambda}\right)^2}{2|x|}}^{1-\frac{a_n|x|}{2}}\left(1-t_n(x)s-\frac{3}{2}a_n|x|t_n(x)s^2\right)^{\alpha}s^{\beta}ds\\
=\;&\int_{\frac{\left(\lambda+\frac{y-x}{2\lambda}\right)^2}{2|x|}}^{1}(1-s)^{\alpha}s^{\beta}ds+\alpha a_n|x|\int_{\frac{\left(\lambda+\frac{y-x}{2\lambda}\right)^2}{2|x|}}^{1}(1-s)^{\alpha-1}s^{\beta+1}\left(1-\frac{3}{2}s\right)ds+o(a_n+c_n)
\end{aligned}$$

holds for large n by arguments similar with (3.15).

With the same arguments as Lemma 3.2, we can get

$$\begin{aligned}
&\frac{1}{2\pi}\int_{\beta_n(t_n(x),t_n(y))}^{\frac{\pi}{2}}\left(1-F\left(t_n(x)/\cos\alpha\right)\right)d\alpha\\
=\;&\frac{1}{4\pi}\int_{\sin^2\beta_n(t_n(x),t_n(y))}^{2a_n|x|-a_n^2x^2}\left(1-F\left(t_n(x)(1-s)^{-\frac{1}{2}}\right)\right)s^{-\frac{1}{2}}(1-s)^{-\frac{1}{2}}ds\\
=\;&\frac{(2a_n|x|)^{\frac{1}{2}}}{4\pi}\left(1-F(t_n(x))\right)\left\{\int_{\frac{\left(\lambda+\frac{y-x}{2\lambda}\right)^2}{2|x|}}^{1}(1-s)^{\alpha}s^{-\frac{1}{2}}ds+A(a_n^{-1})|x|^{-\tau}\int_{\frac{\left(\lambda+\frac{y-x}{2\lambda}\right)^2}{2|x|}}^{1}(1-s)^{\alpha}\frac{(1-s)^{-\tau}-1}{\tau}s^{-\frac{1}{2}}ds\right.
\end{aligned}$$

$$+a_n|x|\int_{\frac{\left(\lambda+\frac{y-x}{2\lambda}\right)^2}{2|x|}}^1 (1-s)^{\alpha-1}s^{\frac{1}{2}}\left(\alpha+1-s-\frac{3}{2}\alpha s\right)ds+o(a_n+c_n+A(a_n^{-1}))$$
$$-\left(\frac{2}{|x|}\right)^{\frac{1}{2}}\left(1-\frac{\left(\lambda+\frac{y-x}{2\lambda}\right)^2}{2|x|}\right)^{\alpha}\left(c_n\gamma\left(1-\frac{y-x}{2\lambda^2}\right)-\frac{a_n}{2}\left(\lambda(y-x)+\frac{x(y-x)}{\lambda}+\frac{3(y-x)^2}{4\lambda}+\frac{(y-x)^3}{8\lambda^3}\right)\right)\Bigg\}$$

for large n, which implies that

$$\begin{aligned}
&\frac{n}{2\pi}\int_{\beta_n(t_n(x),t_n(y))}^{\frac{\pi}{2}}(1-F(t_n(x)/\cos\alpha))\,d\alpha\\
=\;& |x|^{\alpha+\frac{1}{2}}\Bigg\{\frac{1}{c_\alpha}\int_{\frac{\lambda+\frac{y-x}{2\lambda}}{\sqrt{2|x|}}}^1(1-s^2)^\alpha ds+\frac{A(a_n^{-1})}{c_\alpha}\frac{|x|^{-\tau}-1}{\tau}\int_{\frac{\lambda+\frac{y-x}{2\lambda}}{\sqrt{2|x|}}}^1(1-s^2)^\alpha ds\\
&-\frac{a_n}{c_\alpha^2}\int_{-1}^1(1-s^2)^{\alpha-1}s^2\left(\alpha+1-s^2-\frac{3}{2}\alpha s^2\right)ds\int_{\frac{\lambda+\frac{y-x}{2\lambda}}{\sqrt{2|x|}}}^1(1-s^2)^\alpha ds\\
&-\frac{A(a_n^{-1})}{c_\alpha^2}\int_{-1}^1(1-s^2)^\alpha\frac{(1-s^2)^{-\tau}-1}{\tau}ds\int_{\frac{\lambda+\frac{y-x}{2\lambda}}{\sqrt{2|x|}}}^1(1-s^2)^\alpha ds\\
&+\frac{A(a_n^{-1})|x|^{-\tau}}{c_\alpha}\int_{\frac{\lambda+\frac{y-x}{2\lambda}}{\sqrt{2|x|}}}^1(1-s^2)^\alpha\frac{(1-s^2)^{-\tau}-1}{\tau}ds\\
&+\frac{a_n|x|}{c_\alpha}\int_{\frac{\lambda+\frac{y-x}{2\lambda}}{\sqrt{2|x|}}}^1(1-s^2)^{\alpha-1}s^2\left(\alpha+1-s^2-\frac{3}{2}\alpha s^2\right)ds+o(a_n+c_n+A(a_n^{-1}))\\
&-\frac{1}{c_\alpha\sqrt{2|x|}}\left(1-\frac{\left(\lambda+\frac{y-x}{2\lambda}\right)^2}{2|x|}\right)^\alpha\left(c_n\gamma\left(1-\frac{y-x}{2\lambda^2}\right)\right.\\
&\left.-\frac{a_n}{2}\left(\lambda(y-x)+\frac{x(y-x)}{\lambda}+\frac{3(y-x)^2}{4\lambda}+\frac{(y-x)^3}{8\lambda^3}\right)\right)\Bigg\}\\
:=\;& M_{\alpha+\frac{1}{2},\lambda}(x,y). \qquad (4.3)
\end{aligned}$$

Similarly, for the case of $\lambda+\frac{y-x}{2\lambda}<0$ with $\lambda^2+x+y+\frac{(x-y)^2}{4\lambda^2}<0$, we have

$$\begin{aligned}
&\frac{n}{2\pi}\int_{\beta_n(t_n(x),t_n(y))}^{\frac{\pi}{2}}(1-F(t_n(x)/\cos\alpha))\,d\alpha\\
=\;& \frac{n}{4\pi}\int_0^1\left(1-F\left(t_n(x)(1-s)^{-\frac{1}{2}}\right)\right)s^{-\frac{1}{2}}(1-s)^{-\frac{1}{2}}ds\\
&+\frac{n}{4\pi}\int_0^{\sin^2\beta_n(t_n(x),t_n(y))}\left(1-F\left(t_n(x)(1-s)^{-\frac{1}{2}}\right)\right)s^{-\frac{1}{2}}(1-s)^{-\frac{1}{2}}ds\\
=\;& M_{\alpha+\frac{1}{2},\lambda}(x,y)
\end{aligned}$$

for large n, which is consistent with the case of $\lambda+\frac{y-x}{2\lambda}\geq 0$.

Define $\tilde{\beta}_n(t_n(x), t_n(y)) = \tau_n - \beta_n(t_n(x), t_n(y))$. From the second-order condition (1.4) and (4.2), it follows that

$$\begin{aligned}
&\frac{\tilde{\beta}_n(t_n(x), t_n(y))}{\sqrt{a_n}} \\
= \;& \lambda + \frac{x-y}{2\lambda} + \gamma\left(1 - \frac{x-y}{2\lambda^2}\right) c_n \\
&+ \frac{a_n}{2}\left(\frac{\lambda^2}{3} + \frac{\lambda(y-x)}{2} + \frac{x(y-x)}{\lambda} + \frac{(y-x)^2}{2\lambda} + \frac{(y-x)^3}{12\lambda^3}\right) + o(a_n + c_n) \qquad (4.4)
\end{aligned}$$

for large n, which implies that

$$\begin{aligned}
\frac{\sin\tilde{\beta}_n(t_n(x), t_n(y))}{\sqrt{a_n}} = \;& \lambda + \frac{x-y}{2\lambda} + \gamma\left(1 - \frac{x-y}{2\lambda^2}\right) c_n \\
&+ \frac{a_n}{2}\left(\lambda(y-x) + \frac{x(y-x)}{\lambda} + \frac{(y-x)^2}{4\lambda} + \frac{(y-x)^3}{8\lambda^3}\right) \qquad (4.5)
\end{aligned}$$

for large n. Using the similar arguments as above, we can get

$$\begin{aligned}
&\frac{n}{2\pi}\int_{\tilde{\beta}_n(t_n(x),t_n(y))}^{\frac{\pi}{2}} (1 - F(t_n(y)/\cos\alpha))\, d\alpha \\
= \;& |y|^{\alpha+\frac{1}{2}} \left\{ \frac{1}{c_\alpha}\int_{\frac{\lambda+\frac{x-y}{2\lambda}}{\sqrt{2|y|}}}^1 (1-s^2)^\alpha ds + \frac{A(a_n^{-1})}{c_\alpha}\frac{|y|^{-\tau}-1}{\tau}\int_{\frac{\lambda+\frac{x-y}{2\lambda}}{\sqrt{2|y|}}}^1 (1-s^2)^\alpha ds \right. \\
&- \frac{a_n}{c_\alpha^2}\int_{-1}^1 (1-s^2)^{\alpha-1} s^2\left(\alpha + 1 - s^2 - \frac{3}{2}\alpha s^2\right) ds \int_{\frac{\lambda+\frac{x-y}{2\lambda}}{\sqrt{2|y|}}}^1 (1-s^2)^\alpha ds \\
&- \frac{A(a_n^{-1})}{c_\alpha^2}\int_{-1}^1 (1-s^2)^\alpha \frac{(1-s^2)^{-\tau}-1}{\tau} ds \int_{\frac{\lambda+\frac{x-y}{2\lambda}}{\sqrt{2|y|}}}^1 (1-s^2)^\alpha ds \\
&+ \frac{A(a_n^{-1})|y|^{-\tau}}{c_\alpha}\int_{\frac{\lambda+\frac{x-y}{2\lambda}}{\sqrt{2|y|}}}^1 (1-s^2)^\alpha \frac{(1-s^2)^{-\tau}-1}{\tau} ds \\
&+ \frac{a_n|y|}{c_\alpha}\int_{\frac{\lambda+\frac{x-y}{2\lambda}}{\sqrt{2|y|}}}^1 (1-s^2)^{\alpha-1} s^2\left(\alpha + 1 - s^2 - \frac{3}{2}\alpha s^2\right) ds + o(a_n + c_n + A(a_n^{-1})) \\
&- \frac{1}{c_\alpha\sqrt{2|y|}}\left(1 - \frac{\left(\lambda + \frac{x-y}{2\lambda}\right)^2}{2|y|}\right)^\alpha \left(c_n\gamma\left(1 - \frac{x-y}{2\lambda^2}\right)\right. \\
&\left.\left. + \frac{a_n}{2}\left(\lambda(y-x) + \frac{x(y-x)}{\lambda} + \frac{(y-x)^2}{4\lambda} + \frac{(y-x)^3}{8\lambda^3}\right)\right)\right\} \qquad (4.6)
\end{aligned}$$

for large n.

Since $\lambda^2 + x + y + \frac{(x-y)^2}{4\lambda^2} < 0$ ensures that both $\frac{\left(\lambda+\frac{x-y}{2\lambda}\right)^2}{2|y|} < 1$ and $\frac{\left(\lambda+\frac{y-x}{2\lambda}\right)^2}{2|x|} < 1$ hold, and

$$\begin{aligned}\mathbb{P}(\xi_{n1} > t_n(x), \eta_{n1} > t_n(y)) &= \frac{1}{2\pi}\int_{\max(\min(\beta_n(t_n(x),t_n(y)),\frac{\pi}{2}),-\frac{\pi}{2})}^{\frac{\pi}{2}} (1 - F(t_n(x)/\cos\alpha))\, d\alpha \\ &\quad + \frac{1}{2\pi}\int_{\max(\min(\tilde{\beta}_n(t_n(x),t_n(y)),\frac{\pi}{2}),-\frac{\pi}{2})}^{\frac{\pi}{2}} (1 - F(t_n(y)/\cos\alpha))\, d\alpha,\end{aligned}$$

c.f., Hashorva (2006), we have

$$\begin{aligned}&-n\left(1 - \mathbb{P}(\xi_{n1} \le t_n(x), \eta_{n1} \le t_n(y))\right) + |x|^{\alpha+\frac{1}{2}}\psi_\alpha\left(\frac{\lambda+\frac{y-x}{2\lambda}}{\sqrt{2|x|}}\right) + |y|^{\alpha+\frac{1}{2}}\psi_\alpha\left(\frac{\lambda+\frac{x-y}{2\lambda}}{\sqrt{2|y|}}\right) \\ =\ & -n\left(1 - G(t_n(x))\right) + |x|^{\alpha+\frac{1}{2}} - n\left(1 - G(t_n(y))\right) + |y|^{\alpha+\frac{1}{2}} \\ &+\frac{n}{2\pi}\int_{\tilde{\beta}_n(t_n(x),t_n(y))}^{\frac{\pi}{2}} (1 - F(t_n(y)/\cos\alpha))\, d\alpha - |y|^{\alpha+\frac{1}{2}}\left(1 - \psi_\alpha\left(\frac{\lambda+\frac{x-y}{2\lambda}}{\sqrt{2|y|}}\right)\right) \\ &+\frac{n}{2\pi}\int_{\beta_n(t_n(x),t_n(y))}^{\frac{\pi}{2}} (1 - F(t_n(x)/\cos\alpha))\, d\alpha - |x|^{\alpha+\frac{1}{2}}\left(1 - \psi_\alpha\left(\frac{\lambda+\frac{y-x}{2\lambda}}{\sqrt{2|x|}}\right)\right) \\ =\ & Q_{\alpha+\frac{1}{2},\lambda}(x,y)\end{aligned}$$

for large n, where $Q_{\alpha+\frac{1}{2},\lambda}(x,y)$ is the one given by Theorem 2.1. Hence,

$$\begin{aligned}&\mathbb{P}\left(M_{n1} \le t_n(x), M_{n2} \le t_n(y)\right) - H_{\alpha+\frac{1}{2},\lambda}(x,y) \\ =\ & \mathbb{P}^n\left(\xi_{n1} \le t_n(x), \eta_{n1} \le t_n(y)\right) - H_{\alpha+\frac{1}{2},\lambda}(x,y) \\ =\ & H_{\alpha+\frac{1}{2},\lambda}(x,y)\left(\exp\left(n\log\mathbb{P}\left(\xi_{n1} \le t_n(x), \eta_{n1} \le t_n(y)\right) + |x|^{\alpha+\frac{1}{2}}\psi_\alpha\left(\frac{\lambda+\frac{y-x}{2\lambda}}{\sqrt{2|x|}}\right) + |y|^{\alpha+\frac{1}{2}}\psi_\alpha\left(\frac{\lambda+\frac{x-y}{2\lambda}}{\sqrt{2|y|}}\right)\right) - 1\right) \\ =\ & H_{\alpha+\frac{1}{2},\lambda}(x,y)\left(n\log\mathbb{P}(\xi_{n1} \le t_n(x), \eta_{n1} \le t_n(y)) + |x|^{\alpha+\frac{1}{2}}\psi_\alpha\left(\frac{\lambda+\frac{y-x}{2\lambda}}{\sqrt{2|x|}}\right) + |y|^{\alpha+\frac{1}{2}}\psi_\alpha\left(\frac{\lambda+\frac{x-y}{2\lambda}}{\sqrt{2|y|}}\right)\right. \\ &\left.+\frac{1}{2}\left(n\log\mathbb{P}(\xi_{n1} \le t_n(x), \eta_{n1} \le t_n(y)) + |x|^{\alpha+\frac{1}{2}}\psi_\alpha\left(\frac{\lambda+\frac{y-x}{2\lambda}}{\sqrt{2|x|}}\right) + |y|^{\alpha+\frac{1}{2}}\psi_\alpha\left(\frac{\lambda+\frac{x-y}{2\lambda}}{\sqrt{2|y|}}\right)\right)^2(1+o(1))\right) \\ =\ & H_{\alpha+\frac{1}{2},\lambda}(x,y)\left(-n\left(1 - \mathbb{P}(\xi_{n1} \le t_n(x), \eta_{n1} \le t_n(y))\right) + |x|^{\alpha+\frac{1}{2}}\psi_\alpha\left(\frac{\lambda+\frac{y-x}{2\lambda}}{\sqrt{2|x|}}\right) + |y|^{\alpha+\frac{1}{2}}\psi_\alpha\left(\frac{\lambda+\frac{x-y}{2\lambda}}{\sqrt{2|y|}}\right)\right. \\ &-\frac{n}{2}\left(1 - \mathbb{P}(\xi_{n1} \le t_n(x), \eta_{n1} \le t_n(y))\right)^2 - \frac{n}{3}\left(1 - \mathbb{P}(\xi_{n1} \le t_n(x), \eta_{n1} \le t_n(y))\right)^3(1+o(1)) \\ &\left.+\frac{1}{2}\left(n\log\mathbb{P}(\xi_{n1} \le t_n(x), \eta_{n1} \le t_n(y)) + |x|^{\alpha+\frac{1}{2}}\psi_\alpha\left(\frac{\lambda+\frac{y-x}{2\lambda}}{\sqrt{2|x|}}\right) + |y|^{\alpha+\frac{1}{2}}\psi_\alpha\left(\frac{\lambda+\frac{x-y}{2\lambda}}{\sqrt{2|y|}}\right)\right)^2(1+o(1))\right) \\ =\ & H_{\alpha+\frac{1}{2},\lambda}(x,y)\Big(Q_{\alpha+\frac{1}{2},\lambda}(x,y) - \frac{1}{2}n^{-1}\left(|x|^{\alpha+\frac{1}{2}}\psi_\alpha\left(\frac{\lambda+\frac{y-x}{2\lambda}}{\sqrt{2|x|}}\right) + |y|^{\alpha+\frac{1}{2}}\psi_\alpha\left(\frac{\lambda+\frac{x-y}{2\lambda}}{\sqrt{2|y|}}\right)\right)^2 \\ &+o\left(a_n + c_n + A(a_n^{-1}) + n^{-1}\right)\Big)\end{aligned}$$

for large n. $\square$

Because the proofs of Theorem 2.2-2.7 are similar, we only prove Theorem 2.5-2.7 below, and omit the proofs of Theorem 2.2-2.4.

Proof of Theorem 2.5. For large n, we have

$$\begin{aligned}
& \frac{n}{2\pi}\int_{\min\left(\beta_n(t_n(x),t_n(y)),\frac{\pi}{2}\right)}^{\frac{\pi}{2}}\left(1-F(t_n(x)/\cos\alpha)\right)d\alpha \\
= \ & \frac{n}{4\pi}\int_{\min(\sin^2\beta_n(t_n(x),t_n(y)),1)}^{1}\left(1-F(t_n(x)(1-s)^{-\frac{1}{2}})\right)s^{-\frac{1}{2}}(1-s)^{-\frac{1}{2}}ds \\
= \ & \frac{n}{4\pi}(2a_n|x|)^{\frac{1}{2}}\int_{\min\left(\frac{\sin^2\beta_n(t_n(x),t_n(y))}{2a_n|x|},1-\frac{a_n|x|}{2}\right)}^{1-\frac{a_n|x|}{2}}\left(1-F\left(t_n(x)(1-2a_n|x|s)^{-\frac{1}{2}}\right)\right)s^{-\frac{1}{2}}(1-2a_n|x|s)^{-\frac{1}{2}}ds \\
= \ & 0 \qquad (4.7)
\end{aligned}$$

if $\lambda+\frac{y-x}{2\lambda}>\sqrt{2|x|}$; and if $\lambda+\frac{y-x}{2\lambda}<-\sqrt{2|x|}$, by using (3.19) we have

$$\begin{aligned}
& \frac{n}{2\pi}\int_{\max\left(\beta_n(t_n(x),t_n(y)),-\frac{\pi}{2}\right)}^{\frac{\pi}{2}}\left(1-F(t_n(x)/\cos\alpha)\right)d\alpha \\
= \ & \frac{n}{4\pi}\left(\int_0^1\left(1-F\left(t_n(x)(1-s)^{-\frac{1}{2}}\right)\right)s^{-\frac{1}{2}}(1-s)^{-\frac{1}{2}}ds\right. \\
& \left.+\int_0^{\min\left(\sin^2\beta_n(t_n(x),t_n(y)),1\right)}\left(1-F\left(t_n(x)(1-s)^{-\frac{1}{2}}\right)\right)s^{-\frac{1}{2}}(1-s)^{-\frac{1}{2}}ds\right) \\
= \ & \frac{n}{2\pi}(2a_n|x|)^{\frac{1}{2}}\int_0^{1-\frac{a_n|x|}{2}}\left(1-F\left(t_n(x)(1-2a_n|x|s)^{-\frac{1}{2}}\right)\right)s^{-\frac{1}{2}}(1-2a_n|x|s)^{-\frac{1}{2}}ds \\
= \ & |x|^{\frac{1}{2}+\alpha}\left\{1+\frac{a_n(|x|-1)}{c_\alpha}\int_0^1(1-s)^{\alpha-1}s^{\frac{1}{2}}\left(\alpha+1-s-\frac{3}{2}\alpha s\right)ds\right. \\
& \left.+A(a_n^{-1})\left(\frac{|x|^{-\tau}-1}{\tau}+\frac{|x|^{-\tau}-1}{c_\alpha}\int_0^1(1-s)^\alpha\frac{(1-s)^{-\tau}-1}{\tau}s^{-\frac{1}{2}}ds\right)+o\left(a_n+A(a_n^{-1})\right)\right\} \qquad (4.8)
\end{aligned}$$

for large n.

Similarly, for large n we can get

$$\frac{n}{2\pi}\int_{\min\left(\tilde\beta_n(t_n(x),t_n(y)),\frac{\pi}{2}\right)}^{\frac{\pi}{2}}\left(1-F(t_n(y)/\cos\alpha)\right)d\alpha=0 \qquad (4.9)$$

if $\lambda+\frac{x-y}{2\lambda}>\sqrt{2|y|}$; and

$$\frac{n}{2\pi}\int_{\max\left(\tilde\beta_n(t_n(x),t_n(y)),-\frac{\pi}{2}\right)}^{\frac{\pi}{2}}\left(1-F(t_n(y)/\cos\alpha)\right)d\alpha$$

$$
\begin{aligned}
&= |y|^{\frac{1}{2}+\alpha}\left\{1+\frac{a_n(|y|-1)}{c_\alpha}\int_0^1(1-s)^{\alpha-1}s^{\frac{1}{2}}\left(\alpha+1-s-\frac{3}{2}\alpha s\right)ds\right.\\
&\quad \left.+A(a_n^{-1})\left(\frac{|y|^{-\tau}-1}{\tau}+\frac{|y|^{-\tau}-1}{c_\alpha}\int_0^1(1-s)^{\alpha}\frac{(1-s)^{-\tau}-1}{\tau}s^{-\frac{1}{2}}ds\right)+o\left(a_n+A(a_n^{-1})\right)\right\} \quad (4.10)
\end{aligned}
$$

if $\lambda+\frac{x-y}{2\lambda}<-\sqrt{2|y|}$.

Now we only prove case (iii), since the proofs of the rest cases are similar. From (3.19), (4.7) and (4.10), it follows that

$$
\begin{aligned}
&-n\left(1-\mathbb{P}(\xi_{n1}\le t_n(x),\eta_{n1}\le t_n(y))\right)+|x|^{\alpha+\frac{1}{2}}\psi_\alpha\left(\frac{\lambda+\frac{y-x}{2\lambda}}{\sqrt{2|x|}}\right)+|y|^{\alpha+\frac{1}{2}}\psi_\alpha\left(\frac{\lambda+\frac{x-y}{2\lambda}}{\sqrt{2|y|}}\right)\\
=\;& -n\left(1-G(t_n(x))\right)+|x|^{\alpha+\frac{1}{2}}-n\left(1-G(t_n(y))\right)+|y|^{\alpha+\frac{1}{2}}\\
&+\frac{n}{2\pi}\int_{\max\left(\tilde{\beta}_n(t_n(x),t_n(y)),-\frac{\pi}{2}\right)}^{\frac{\pi}{2}}\left(1-F(t_n(y)/\cos\alpha)\right)d\alpha-|y|^{\alpha+\frac{1}{2}}\\
&+\frac{n}{2\pi}\int_{\min\left(\beta_n(t_n(x),t_n(y)),\frac{\pi}{2}\right)}^{\frac{\pi}{2}}\left(1-F(t_n(x)/\cos\alpha)\right)d\alpha\\
=\;& -|x|^{\frac{1}{2}+\alpha}\left(\frac{a_n(|x|-1)}{c_\alpha}\int_0^1(1-s)^{\alpha-1}s^{\frac{1}{2}}\left(\alpha+1-s-\frac{3}{2}\alpha s\right)ds\right.\\
&\left.+A(a_n^{-1})\left(\frac{|x|^{-\tau}-1}{\tau}+\frac{|x|^{-\tau}-1}{c_\alpha}\int_0^1(1-s)^{\alpha}\frac{(1-s)^{-\tau}-1}{\tau}s^{-\frac{1}{2}}ds\right)+o\left(a_n+A(a_n^{-1})\right)\right)
\end{aligned}
$$

for large n, then (2.6) holds, which complete the proof. □

Proof of Theorem 2.6. (i) For the case of $\rho_n\equiv 1$. Note that

$$
\mathbb{P}\left(M_{n1}\le 1+a_nx,M_{n2}\le 1+a_ny\right)=G^n\left(1+a_n\min(x,y)\right)
$$

and

$$
H_{\alpha,0}(x,y)=\Psi_{\alpha+\frac{1}{2}}\left(\min(x,y)\right)=\exp\left(-|\min(x,y)|^{\alpha+\frac{1}{2}}\right).
$$

It follows from (3.19) that (2.8) holds.

(ii) If $\rho_n\in(0,1)$ such that $\frac{a_n^{5/3}}{1-\rho_n}\to 0$ as $n\to\infty$, which implies that

$$
\frac{\frac{t_n(y)}{t_n(x)}-\rho_n}{\sqrt{a_n}\sin\tau_n}=\lambda_n+\frac{y-x}{2\lambda_n}-\frac{(y-x)x}{2\lambda_n}a_n+o\left(\frac{a_n}{\lambda_n}\right).
$$

Hence,

$$\frac{\sin\beta_n(t_n(x),t_n(y))}{\sqrt{a_n}}=\lambda_n+\frac{y-x}{2\lambda_n}-\frac{(y-x)^3a_n}{16\lambda_n^3}+o\left(\frac{a_n}{\lambda_n^3}\right).$$

Similarly,

$$\frac{\sin\tilde{\beta}_n(t_n(x),t_n(y))}{\sqrt{a_n}}=\lambda_n+\frac{x-y}{2\lambda_n}+\frac{(y-x)^3a_n}{16\lambda_n^3}+o\left(\frac{a_n}{\lambda_n^3}\right)$$

for large n. Hence $\beta_n\left(t_n(\min(x,y)),t_n(\max(x,y))\right)>0$ and $\tilde{\beta}_n\left(t_n(\min(x,y)),t_n(\max(x,y))\right)<0$ for large n. Using (3.19), for large n we have

$$\begin{aligned}
&\frac{n}{2\pi}\int_{\min\left(\beta_n(t_n(\min(x,y)),t_n(\max(x,y))),\frac{\pi}{2}\right)}^{\frac{\pi}{2}}\left(1-F(t_n(\min(x,y))/\cos\alpha)\right)d\alpha\\
=\;&\frac{n(2a_n|\min(x,y)|)^{\frac{1}{2}}}{4\pi}\int_{\min\left(\frac{\sin^2\beta_n(t_n(\min(x,y)),t_n(\max(x,y)))}{2a_n|\min(x,y)|},1-\frac{a_n|\min(x,y)|}{2}\right)}^{1-\frac{a_n|\min(x,y)|}{2}}s^{-\frac{1}{2}}(1-2a_n|\min(x,y)|s)^{-\frac{1}{2}}\\
&\times\left(1-F\left(t_n(\min(x,y))(1-2a_n|\min(x,y)|s)^{-\frac{1}{2}}\right)\right)ds\\
=\;&0
\end{aligned}\tag{4.11}$$

and

$$\begin{aligned}
&\frac{n}{2\pi}\int_{\max\left(\tilde{\beta}_n(t_n(\min(x,y)),t_n(\max(x,y))),-\frac{\pi}{2}\right)}^{\frac{\pi}{2}}\left(1-F(t_n(\max(x,y))/\cos\alpha)\right)d\alpha\\
=\;&\frac{n}{2\pi}\int_0^{\frac{\pi}{2}}\left(1-F(t_n(\max(x,y))/\cos\alpha)\right)d\alpha\\
&+\frac{n}{2\pi}\int_0^{\min\left(-\tilde{\beta}_n(t_n(\min(x,y)),t_n(\max(x,y))),\frac{\pi}{2}\right)}\left(1-F(t_n(\max(x,y))/\cos\alpha)\right)d\alpha\\
=\;&|\max(x,y)|^{\frac{1}{2}+\alpha}\left\{1+\frac{a_n(|\max(x,y)|-1)}{c_\alpha}\int_0^1(1-s)^{\alpha-1}s^{\frac{1}{2}}\left(\alpha+1-s-\frac{3}{2}\alpha s\right)ds\right.\\
&+A(a_n^{-1})\left(\frac{|\max(x,y)|^{-\tau}-1}{c_\alpha}\int_0^1(1-s)^\alpha\frac{(1-s)^{-\tau}-1}{\tau}s^{-\frac{1}{2}}ds\right.\\
&\left.\left.+\frac{|\max(x,y)|^{-\tau}-1}{\tau}\right)+o\left(a_n+A(a_n^{-1})\right)\right\}.
\end{aligned}\tag{4.12}$$

Combining (4.11) and (4.12), we have

$$\begin{aligned}
& n\Big(1-\mathbb{P}(\xi_{n1}\le t_n(\min(x,y)),\eta_{n1}\le t_n(\max(x,y)))\Big)-|\min(x,y)|^{\alpha+\frac{1}{2}} \\
= \;& |\min(x,y)|^{\frac{1}{2}+\alpha}\left\{\frac{a_n(|\min(x,y)|-1)}{c_\alpha}\int_0^1(1-s)^{\alpha-1}s^{\frac{1}{2}}\left(\alpha+1-s-\frac{3}{2}\alpha s\right)ds\right. \\
& +A(a_n^{-1})\left(\frac{|\min(x,y)|^{-\tau}-1}{c_\alpha}\int_0^1(1-s)^{\alpha}\frac{(1-s)^{-\tau}-1}{\tau}s^{-\frac{1}{2}}ds\right. \\
& \left.\left.+\frac{|\min(x,y)|^{-\tau}-1}{\tau}\right)+o\left(a_n+A(a_n^{-1})\right)\right\}
\end{aligned}$$

for large n, which implies that (2.6) also holds. The proof is complete. □

Proof of Theorem 2.7. (i) By $\lim_{n\to\infty}\frac{1-\rho_n}{a_n}=\infty$ and $\lim_{n\to\infty}\frac{(1-\rho_n)^3}{a_n}=0$, we have

$$\frac{\frac{t_n(y)}{t_n(x)}-\rho_n}{\sqrt{a_n}\sin\tau_n}=\lambda_n+\frac{y-x}{2\lambda_n}+\frac{a_n\lambda_n^3}{2}+o(a_n\lambda_n^3)$$

for large n, which implies that

$$\frac{\sin\beta_n(t_n(x),t_n(y))}{\sqrt{a_n}}=\lambda_n+\frac{y-x}{2\lambda_n}+o(a_n\lambda_n^3) \tag{4.13}$$

and

$$\frac{\sin\tilde{\beta}_n(t_n(x),t_n(y))}{\sqrt{a_n}}=\lambda_n+\frac{x-y}{2\lambda_n}+o(a_n\lambda_n^3). \tag{4.14}$$

Here, (4.13) and (4.14) show that $\beta_n(t_n(x),t_n(y))>0$ and $\tilde{\beta}_n(t_n(x),t_n(y))>0$ for large n. Then combining with (3.19),

$$\begin{aligned}
& -n\,\mathbb{P}(X\le t_n(x),Y\le t_n(y))+|x|^{\alpha+\frac{1}{2}}+|y|^{\alpha+\frac{1}{2}} \\
= \;& -n(1-G(t_n(x)))+|x|^{\alpha+\frac{1}{2}}-n(1-G(t_n(y)))+|y|^{\alpha+\frac{1}{2}} \\
= \;& -|x|^{\frac{1}{2}+\alpha}\left(\frac{a_n(|x|-1)}{c_\alpha}\int_0^1(1-s)^{\alpha-1}s^{\frac{1}{2}}\left(\alpha+1-s-\frac{3}{2}\alpha s\right)ds\right. \\
& \left.+A(a_n^{-1})\left(\frac{|x|^{-\tau}-1}{\tau}+\frac{|x|^{-\tau}-1}{c_\alpha}\int_0^1(1-s)^{\alpha}\frac{(1-s)^{-\tau}-1}{\tau}s^{-\frac{1}{2}}ds\right)\right) \\
& -|y|^{\frac{1}{2}+\alpha}\left(\frac{a_n(|y|-1)}{c_\alpha}\int_0^1(1-s)^{\alpha-1}s^{\frac{1}{2}}\left(\alpha+1-s-\frac{3}{2}\alpha s\right)ds\right.
\end{aligned}$$

$$+A(a_n^{-1})\left(\frac{|y|^{-\tau}-1}{\tau}+\frac{|y|^{-\tau}-1}{c_\alpha}\int_0^1(1-s)^\alpha\frac{(1-s)^{-\tau}-1}{\tau}s^{-\frac{1}{2}}ds\right)+o\left(a_n+A(a_n^{-1})\right)$$

for large n. Hence we can derive (2.2) with $\lambda=\infty$.

(ii) For $\lim_{n\to\infty}\rho_n=d\in[-1,1)$, we have

$$\beta_n(t_n(x),t_n(y))\to\arctan\frac{1-d}{\sin(\arccos d)}=\frac{1}{2}\arccos d>0$$

and

$$\tilde{\beta}_n(t_n(x),t_n(y))\to\arccos d-\arctan\frac{1-d}{\sin(\arccos d)}=\frac{1}{2}\arccos d>0$$

as $n\to\infty$. Using the same arguments as the case of (i), the desired result is derived, which complete the proof. □

Acknowledgements This work was supported by the National Natural Science Foundation of China (grant No. 11171275 and No. 11501113), the Key Project of Fujian Education Committee (grant No. JA15045), the Hujiang Fund for Ph.D. Scientific Research Startup (grant No. BSQD201608).

References

[1] de Haan, L. and Ferreira, A. (2006). *Extreme Value Theory: An Introduction*. Springer-verlag, Berlin.

[2] Frick, M. and Reiss, R-D. (2010). Limiting distributions of maxima under triangular schemes. *Journal of Multivariate Analysis*, **10**, 2563–2568.

[3] Frick, M. and Reiss, R-D. (2013). Expansions and penultimate distributions of maxima of bivariate Gaussian random vectors. *Statistics and Probability Letters*, **83**, 2563–2568.

[4] Hashorva, E. (2005). Elliptical triangular arrays in the max-domain of attraction of Hüsler-Reiss distribution. *Statistics and Probability Letters*, **72**, 125–135.

[5] Hashorva, E. (2006). On the max-domain of attractions of bivariate elliptical arrays. *Extremes*, **8**, 225–233.

[6] Hashorva, E. (2008). Extremes of weighted dirichlet arrays. *Extremes*, **4**, 393–420.

[7] Hashorva, E. and Ling, C. (2016). Maxima of skew elliptical triangular arrays, *Communications in Statistics-Theory and Methods*, **12**, 3692–3705.

[8] Hashorva, E., Kabluchko, Z., and Wübker, A. (2012). Extremes of independent chi-square random vectors. *Extremes*, **15**, 35–42.

[9] Hashorva, E., Peng, L. and Weng, Z. (2015). Maxima of a triangular array of multivariate Gaussian sequence. *Statistics and Probability Letters*, **103**, 62–72.

[10] Hashorva, E., Peng, Z. and Weng, Z. (2016). Higher-order expansions of distributions of maxima in a Hüsler-Reiss model. *Methodology and Computing in Applied Probability*, **18**, 181–196.

[11] Hashorva, E. and Weng, Z. (2013). Limit laws for extremes of dependent stationary Gaussian arrays. *Statistics and Probability Letters*, **83**, 320-330.

[12] Hüsler, J. and Reiss, R-D. (1989). Maxima of normal random vectors: Between independence and complete dependence. *Statistics and Probability Letters*, **7**, 283–286.

[13] Kabluchko, Z., de Haan, L. and Schlatter, M. (2009). Stationary max-stable fields associated to negative definite functions. *The Annals of Probability*, **37**, 2042–2065.

[14] Liao X. and Peng Z. (2014). Convergence rate of maxima of bivariate Gaussian arrays to the Hüsler-Reiss distribution. *Statistics and Its Interface*, **7(3)**, 351–362.

[15] Liao X. and Peng Z. (2015). Asymptotics for the maxima and minima of Hüsler-Reiss bivariate Gaussian arrays. *Extremes*, **18(1)**, 1–14.

[16] Resnick, S.I. (1987). *Extreme Values, Regular Variation and Point Processes*. New York: Springer verlag.